Gastrointestinal Associates of Rockland 500 New Hempstead Rd New City NY 10956 845-362-3200

Fax: 845-290-8180

NULYTELY OR TRILYTE /DULCOLAX ACCELERATED COMBO PREPARATION FOR COLONOSCOPY

*Unless directed by the physician, if you take heart or blood pressure medication, you may take with only a sip of water at your normally scheduled time as well as take your inhaler at your normally scheduled time. *If you are taking Warfarin, Coumadin, Plavix, clopidogrel bisulfate, or diabetic medication, discuss this with the physician prior to your scheduled procedure date.

- STOP PRADAXA FIVE (5) DAYS PRIOR TO PROCEDURE.
- STOP EFFIENT SEVEN (7) DAYS PRIOR TO PROCEDURE.
- No aspirin or aspirin containing products, Alka-seltzer, anti-arthritis or anti-inflammatory medications for seven (7) days prior to the scheduled procedure date. Tylenol is fine.
- No Metamucil, Fibercon, Citrucel or iron supplements five (5) days before your procedure.
- No vegetables for two (2) days before the procedure.

Fill your prescription for Nulytely or Trilyte and purchase Dulcolax laxatives over the counter.

Please follow the preparation carefully and call the office if you have any questions:

24 HOURS BEFORE YOUR PROCEDURE:

Start a clear liquid d	iet – water, apple juice, o	coffee, tea (without milk or nondairy creamer), broth,
Jello, ices, (nothing	red or purple).	
AT	M THE	YOUR PROCEDURE
Take the 4 Dulcolax	laxatives. Follow with a	another 8 oz of approved clear liquid and continue with
the clear liquid die	t until you begin the Nu	ılytely or Trilyte.
A T	M THE	YOUR PROCEDURE
Following the instru	ctions for mixture of the	Nulytely or Trilyte, you will prepare 4 liters of the
liquid. Refrigerate	prior to drinking for be	etter taste ONLY DRINK 2 LITERS OF THE
MIXTURE AND F	INISH THE 2 LITERS	BBYM. DISCARD THE REMAINDER OF
THE MIXTURE.	You may continue clear	liquids until five hours prior to your procedure and
then nothing else to	o drink until after your	procedure time. Please bring your medication list with
vou!	•	•

You must have someone to drive you home following the procedure due to the medication given at the time of the procedure. Please check with your insurance regarding pre-certification.