



ADVANCED SINUS
AND
ALLERGY CENTER

1030 W. Higgins #325
Park Ridge, IL 60068

**Post-Operative Instructions after Septoplasty, Inferior Turbinate Reduction,
Latera Implantation and Uvulectomy**

Diet:

It is important that you drink plenty of fluids to prevent dehydration and to help soothe the throat pain

No crunchy foods like chips or pizza, no acidic foods or very hot liquids allowed for 2 weeks

No straws

Liquid diet only for the first 3 days: fruit juice, milkshakes, ice cream, popsicles, pudding, Carnation breakfast drink, Gatorade, Gingerale and Pedialyte

Soft diet only from day 4 to 14: mashed fruits, oatmeal, soft cooked pasta noodles, scrambled eggs, mashed potatoes, jello, and anything from the list above

Pain:

It is common to have a sore throat, ear pain, nose pain and pain while opening your jaw after this surgery. This will get better every day.

Take your pain medication (Lortab) every 4 hours while awake for the first 3 days.

After 3 days you may alternate Lortab and Tylenol liquid (which you can buy over the counter) but DO NOT take both together because they can cause liver toxicity since they both contain Acetaminophen

Do not take more than 4 grams of Tylenol (Acetaminophen) per 24 hours

Do not drive while taking Lortab because it is a narcotic medication

Take milk of magnesia daily while you are taking Lortab to prevent constipation

DO NOT TAKE any Aspirin or Aspirin-containing products like Motrin, Ibuprofen, Nuprin, Aleve, Celebrex, Naproxyn, Excedrin, Advil or Peptobismol for 2 weeks after surgery

Migraine medications should be used only after discussion with your physician and should not be taken at the same time as your narcotic pain medication

Nasal Care:



ADVANCED SINUS
AND
ALLERGY CENTER

1030 W. Higgins #325
Park Ridge, IL 60068

Change the mustache dressing as needed. The drainage from your nose will slow down and will stop after 24-48 hours

Clean the front of the nose with a Qtip dipped in hydrogen peroxide three times per day and apply Bacitracin or Neosporin ointment to the front of the nose with a clean Qtip after cleansing

Do not pull at the splints or the blue suture inside your nose

Do not squint your nose or remove the white bandaids over the nose. Do not wear glasses for one week after surgery. Wash your face with a cloth and do not get the white bandaids wet.

Use a humidifier or vaporizer in your room during the night to prevent throat pain with mouth breathing

During the first 7 days while the splints are in place, spray 3 sprays of nasal saline (salt water spray) in each nostril every 2 hours while awake. After spraying, snort back the saline and mucous and swallow it or spit it out, but do not blow your nose

The day after the splints are removed, start rinsing your nose with saline using a NeilMed bottle 3 times daily. You can purchase the bottle, the distilled water and the salt Neil Med packets at your local pharmacy.

This will keep the inside of the splints clean and allow you to breathe better through your nose

If you develop bright red blood from your nose:

1. Spray 4 sprays of Afrin (Oxymetazoline) in each side of your nose and pinch the front of the nose for 15 minutes using a nose clip AND
2. Apply an ice pack or a bag of frozen peas over the outside of the nose AND
3. Suck on ice chips
4. If the bleeding does not stop after 15 minutes, call our office. If it is after hours, follow the instructions on the phone and have the provider on call paged.

Activities:

No nose blowing for 2 weeks after surgery

Cough and sneeze with your mouth open for 2 weeks after surgery

No heavy lifting, bending or straining for 2 weeks after surgery

No strenuous exercise for 2 weeks after surgery

Sleep with the head of bed elevated with extra pillows or in a recliner if you prefer



ADVANCED SINUS
AND
ALLERGY CENTER

1030 W. Higgins #325
Park Ridge, IL 60068

You can begin showering the day after surgery. Steam showers are recommended. Do not bend your head while showering.

When to call your doctor:

Pain that does not respond to the pain medication

Temperature over 103F

Persistent cough

Coughing/spitting up blood or profuse bleeding from your nose

Nausea and vomiting

Dehydration (decrease urination or when your urine turns very dark yellow or amber)

Yellow/green drainage from the nose and foul odor in the nose

Other restrictions:

You cannot travel by airplane for 2 weeks after surgery

You cannot travel outside of the Chicago area for 2 weeks after surgery

No smoking or chewing tobacco or drinking alcohol for 2 weeks after surgery

No smoking marijuana or using other illicit drugs

What to buy at the pharmacy

Prescribed medications: antibiotic (liquid Augmentin) and Lortab or Hycet (pain medication)

Over-the-counter medications: nasal saline (Ocean Spray), Afrin (Oxymetazoline), Milk of Magnesia and liquid Tylenol (Acetaminophen)

Other supplies: nose clip, a box of small gauze and a roll of skin tape, a Neil Med bottle, Neil Med salt packets and distilled water

How to get in touch with your doctor:

1. During regular business hours, please call our office at (847) 655-7442
2. For emergencies after hours, please call the same number and follow the prompts to have the provider on call paged
3. For non-emergencies after hours, you may leave a message on the machine and we will return your call the next morning



ADVANCED SINUS
AND
ALLERGY CENTER

1030 W. Higgins #325
Park Ridge, IL 60068