## SURGERY DISCHARGE INSTRUCTIONS

Patient:	ent: Date:					
1.	Have prescriptions filled immediately. Take prescriptions ONLY as directed:  Durcicef/Erythromycin 500mg twice daily and Percocet/Vicodin 5mg every 4 hours as needed Additional:					
	could cause you to feel faint. Take medication with food.					
2.	For the first 72 hours 93) days following surgery, REMAIN OFF OF YOUR FEET as much as possible. Elevate your foot above heart level with your knee slightly bent. Limit activity since over activity causes swelling and pain.					
3.	WALKING INSTRUCTIONS:  No weight on operated foot					
	Walk only with surgical shoe or removable cast. Walking must be kept to a minimum for 3 days. Additional Instructions:					
4.	Do NOT remove dressings or allow them to get wet. Cast/bandage shower covers are available for purchasin our office.					
5.	Put ice on surgical area, 30 min per hour while awake, this will help with swelling and pain.					
6.	Keep pets AWAY from dressing to prevent infection.					
7.	CALL THE OFFICE IMMEDIATELY IF:					
	If excessive bleeding occurs					
C)	Bandage/cast feels too tight causing numbness and tingling Bandage/cast gets wet					
	Fever or chills occur					
F)	Toes become white or bluish -It is very important that you monitor the color and temperature of the toes. The capillary refill should return instantly to all digits including on the toe that was operated on. The temperature of the toe should be warm. The color should be pink. If not please call Dr. Polun immediately.					
8.	If your left foot is in a surgical shoe/cast, you may drive vehicles equipped with automatic drive. If your right foot is in a surgical shoe/cast you may NOT drive for 3 days.					
9.	We recommend Mederma be applied to the incision site after sutures and cast are removed to avoid scarring.					
POST C	OPERATIVE APPOINTMENT:					
Patient S	Signature Physician Signature					

Dr. Franklin R. Polun Office: 301-983-8202 Cell 301-529-1575