

Stroke

Definition

A stroke is an interruption of the blood supply to any part of the brain. A stroke is sometimes called a "brain attack."

See also:

- Aneurysm
- Arteriovenous malformation (AVM)
- Cerebral hemorrhage

Alternative Names

Cerebrovascular disease; CVA; Cerebral infarction; Cerebral hemorrhage; Ischemic stroke; Stroke - ischemic; Cerebrovascular accident; Stroke - hemorrhagic

Causes

A stroke happens when blood flow to a part of the brain is interrupted because a blood vessel in the brain is blocked or bursts open.

If blood flow is stopped for longer than a few seconds, the brain cannot get blood and oxygen. Brain cells can die, causing permanent damage.

There are two major types of stroke: ischemic stroke and hemorrhagic stroke.



ISCHEMIC STROKE

Ischemic stroke occurs when a blood vessel that supplies blood to the brain is blocked by a blood clot. This may happen in two ways:

- A clot may form in an artery that is already very narrow. This is called a thrombus. If it completely blocks the artery, it is called a *thrombotic stroke*.
- A clot may break off from another place in the blood vessels of the brain, or some other part of the body, and travel up to the brain to block a smaller artery. This is called an embolism. It causes an *embolic stroke*.

Ischemic strokes may result from clogged arteries, a condition called atherosclerosis. This may affect the arteries within the brain or the arteries in the neck that carry blood to the brain. Fat, cholesterol, and other substances collect on the wall of the arteries, forming a sticky substance called plaque. Over time, the plaque builds up. This often makes it hard for blood to flow properly, which can cause the blood to clot.

Ischemic strokes may also be caused by blood clots that form in the heart or other parts of the body. These clots travel through the blood and can get stuck in the small arteries of the brain. This is known as a cerebral embolism.



Certain drugs and medical conditions can make your blood more likely to clot and raise your risk for ischemic stroke. A common cause of ischemic stroke in people under age 40 is carotid dissection, or a tear in the lining of the carotid artery. The tear lets blood flow between the layers of the carotid artery. This causes narrowing of the carotid artery that is not due to plaque buildup.

Some ischemic strokes start out without any bleeding, and then bleeding into the damaged area occurs.

HEMORRHAGIC STROKE

Hemorrhagic stroke occurs when a blood vessel in part of the brain becomes weak and bursts open, causing blood to leak into the brain. Some people have defects in the blood vessels of the brain that make this more likely. The flow of blood that occurs after the blood vessel ruptures damages brain cells.

STROKE RISKS

High blood pressure is the number one risk factor for strokes. The following also increase your risk for stroke:

- Atrial fibrillation
- Diabetes
- Family history of stroke
- Heart disease
- High cholesterol
- Increasing age

Certain medications make blood clots more likely, and therefore increase your chances for a stroke. Birth control pills can increase the chances of having blood clots, especially in woman who smoke and who are older than 35.

Men have more strokes than women. However, women have a higher risk of stroke during pregnancy and in the weeks immediately after pregnancy.

The following can increase the risk of bleeding into the brain, which makes you more likely to have a stroke:

- Alcohol use
- Bleeding disorders
- Cocaine use
- Head injury

For more information see: [Stroke risk factors and prevention](#)

Symptoms

The symptoms of stroke depend on what part of the brain is damaged. In some cases, a person may not even be aware that he or she has had a stroke.

Symptoms usually develop suddenly and without warning, or they may occur on and off for the first day or two. Symptoms are usually most severe when the stroke first happens, but they may slowly get worse.

A headache may occur, especially if the stroke is caused by bleeding in the brain. The headache:

- Starts suddenly and may be severe
- Occurs when lying flat
- Wakes you up from sleep
- Gets worse when you change positions or when you bend, strain, or cough

Other symptoms depend on the severity of the stroke and what part of the brain is affected. Symptoms may include:

- Change in alertness (including sleepiness, unconsciousness, and coma)
- Changes in hearing
- Changes in taste
- Clumsiness
- Confusion or loss of memory
- Difficulty swallowing
- Difficulty writing or reading
- Dizziness or abnormal sensation of movement (vertigo)
- Lack of control over the bladder or bowels
- Loss of balance
- Loss of coordination
- Muscle weakness in the face, arm, or leg (usually just on one side)

- Numbness or tingling on one side of the body
- Personality, mood, or emotional changes
- Problems with eyesight, including decreased vision, double vision, or total loss of vision
- Sensation changes that affect touch and the ability to feel pain, pressure, different temperatures, or other stimuli
- Trouble speaking or understanding others who are speaking
- Trouble walking

Exams and Tests

A complete physical and neurological exam should be performed. Your doctor will:

- Check for problems with vision, movement, sensation, reflexes, understanding, and speaking. Your doctor and nurses will repeat this exam over time to see if your stroke is getting worse or is improving.
- Listen for an abnormal sound, called a "bruit," when using a stethoscope to listen to the carotid arteries in the neck. A bruit is caused by turbulent blood flow.
- Check and assess your blood pressure, which may be high.

Tests can help your doctor determine the type, location, and cause of the stroke and rule out other disorders that may be responsible for the symptoms.

- A CT scan of the brain is often done soon after symptoms of a stroke begin. An MRI scan of the brain may be done instead or afterwards.
- Magnetic resonance angiography (MRA) or CT angiography may be done to check for abnormal blood vessels in the brain that may have caused the stroke.
- Echocardiogram may be done if the stroke could have been caused by a blood clot from the heart.
- Carotid duplex (a type of ultrasound exam) can show if narrowing of the neck arteries (carotid stenosis) led to the stroke.
- An angiogram of the head can reveal which blood vessel is blocked or bleeding, and help your doctor decide if the artery can be reopened using a thin tube.
- Laboratory tests will include a complete blood count (CBC), bleeding time, and blood clotting tests (prothrombin time or partial thromboplastin time). They will also check your blood cholesterol and sugar.
- Electrocardiogram (ECG) and heart rhythm monitoring can help determine if an irregular heartbeat (such as atrial fibrillation) caused the stroke.
- A spinal tap (cerebrospinal fluid exam) may also be done.

Treatment

A stroke is a medical emergency. Immediate treatment can save lives and reduce disability. Call 911 or your local emergency number or seek immediate medical care at the first signs of a stroke.

It is very important for people who are having stroke symptoms to get to a hospital as quickly as possible. If the stroke is caused by a blood clot, a clot-busting drug may be given to dissolve the clot.

Most of the time, patients must reach a hospital within 3 hours after symptoms begin. Some people may be able to receive these drugs for up to 4 - 5 hours after symptoms begin.

Treatment depends on the severity and cause of the stroke. A hospital stay is required for most strokes.

TREATMENT IN THE HOSPITAL

Clot-busting drugs (thrombolytic therapy) may be used if the stroke is caused by a blood clot. Such medicine breaks up blood clots and helps restore blood flow to the damaged area. However, not everyone can receive this type of medicine.

- For these drugs to work, a person must be seen and treatment must begin within 3 hours of when the symptoms first started. A CT scan must be done to see whether the stroke is from a clot or from bleeding.
- If the stroke is caused by bleeding rather than clotting, clot-busting drugs (thrombolytics) can cause more bleeding.

Other treatments depend on the cause of the stroke:

- Blood thinners such as heparin or warfarin (Coumadin) may be used to treat strokes due to blood clots. Aspirin or clopidogrel (Plavix) may also be used.
- Other medications may be needed to control other symptoms, including high blood pressure. Painkillers may be given to control severe headache.
- In some situations, a special stroke team and skilled radiologists may be able to use angiography to highlight the clogged blood vessel and open it up.

- For hemorrhagic stroke, surgery is often required to remove blood from around the brain and to repair damaged blood vessels.
- Surgery on the carotid artery may be needed.

See also:

- Carotid artery disease
- Carotid artery surgery

Nutrients and fluids may be necessary, especially if the person has swallowing difficulties. These may be given through a vein (intravenously) or a feeding tube in the stomach (gastrostomy tube). Swallowing difficulties may be temporary or permanent.

Physical therapy, occupational therapy, speech therapy, and swallowing therapy will all begin in the hospital.

LONG-TERM TREATMENT

The goal of treatment after a stroke is to help the patient recover as much function as possible and prevent future strokes.

The recovery time and need for long-term treatment differs from person to person. Problems moving, thinking, and talking often improve in the weeks to months after a stroke. A number of people who have had a stroke will still continue to improve in the months or years after the stroke.

See: Stroke recovery for information about:

- Bladder and bowel problems
- Living at home rather than a nursing facility
- Muscle and nerve problems
- Speech problems
- Stroke rehabilitation
- Swallowing and eating problems
- Thinking and memory problems

Support Groups

Additional support and resources are available from the American Stroke Association -- www.strokeassociation.org.

Outlook (Prognosis)

The outlook depends on the type of stroke, how much brain tissue is damaged, what body functions have been affected, and how quickly treatment is received. Recovery may occur completely, or there may be some permanent loss of function.

Over half of the people who have a stroke are able to function independently at home.

If treatment with clot-busting drugs is successful, the symptoms of a stroke may completely go away. However, patients do not often arrive at the hospital soon enough to receive these drugs, or there are complicating medical conditions that prevent their use.

People who have an ischemic stroke (stroke due to a blood clot) have a better chance of surviving than those who have a hemorrhagic stroke (stroke due to bleeding in the brain).

The risk for a second stroke is highest over the first few weeks or months after the first stroke and then begins to lessen.

Possible Complications

- Breathing food into the airway (aspiration)
- Decreased life span
- Difficulty communicating
- Fractures
- Malnutrition
- Muscle spasticity
- Permanent loss of brain functions
- Permanent loss of movement or sensation in one or more parts of the body
- Problems due to loss of mobility, including joint contractures and pressure sores

- Reduced ability to function or care for self
- Reduced social interactions
- Side effects of medications

When to Contact a Medical Professional

Stroke is a medical emergency that requires immediate treatment. Call your local emergency number (such as 911) if someone has symptoms of a stroke.

Prevention

To help prevent a stroke:

- Avoid fatty foods. Follow a healthy, low-fat diet.
- Do not drink more than 1 to 2 alcoholic drinks a day.
- Exercise regularly: 30 minutes a day if you are not overweight; 60 - 90 minutes a day if you are overweight.
- Get your blood pressure checked every 1 - 2 years, especially if high blood pressure runs in your family.
- Have your cholesterol checked. If you are at high risk for stroke, your LDL "bad" cholesterol should be lower than 100 mg/dL. Your doctor may recommend that you try to reduce your LDL cholesterol to 70 mg/dL.
- Follow your doctor's treatment recommendations if you have high blood pressure, diabetes, high cholesterol, and heart disease.
- Quit smoking.

Aspirin therapy (81 mg a day or 100 mg every other day) is recommended for stroke prevention in all men who have stroke risk factors, and in women under age 65 who are at risk for stroke, as long as the benefits outweigh the risks. It should be considered for women over age 65 only if their blood pressure is controlled and the benefit is greater than the risk of gastrointestinal bleeding and brain hemorrhage. Ask your doctor if aspirin is right for you.

Your doctor may also recommend aspirin therapy or another blood thinner if you have had a transient ischemic attack (TIA) or stroke in the past, or if you currently have:

- Congestive heart failure
- Irregular heartbeat (such as atrial fibrillation)
- Mechanical heart valve
- Other risk factors for stroke

A type of surgery called carotid endarterectomy may help prevent new strokes from occurring in persons with large blockages in their neck arteries.

See also: Stroke risk factors and prevention

References

Latchaw RE, Alberts MJ, Lev MH, Connors JJ, Harbaugh RE, Higashida RT, et al. Recommendations for imaging of acute ischemic stroke: a scientific statement from the American Heart Association. *Stroke*. 2009;40:3646-3678. Epub 2009 Sep 24.

Del Zoppo GJ, Saver JL, Jauch EC, Adams HP Jr: American Heart Association Stroke Council, Expansion of the time window for treatment of acute ischemic stroke with intravenous tissue plasminogen activator: a science advisory from the American Heart Association/American Stroke Association. *Stroke*. 2009;40:2945-2948. Epub 2009 May 28.

Adams RJ, Albers G, Alberts MJ, Benavente O, Furie K, Goldstein LB, et al. Update to the AHA/ASA recommendations for the prevention of stroke in patients with stroke and transient ischemic attack. *Stroke*. 2008;39:1647-1652. Epub 2008 Mar 5.

Chung CS, Caplan LR. Stroke and other neurovascular disorders. In: Goetz, CG, eds. *Textbook of Clinical Neurology*. 3rd ed. Philadelphia, Pa: Saunders Elsevier; 2007:chap 45.



A.D.A.M., Inc. is accredited by URAC, also known as the American Accreditation HealthCare Commission (www.urac.org). URAC's accreditation program is an independent audit to verify that A.D.A.M. follows rigorous standards of quality and accountability. A.D.A.M. is among the first to achieve this important distinction for online health information and services. Learn more about A.D.A.M.'s editorial policy, editorial process and privacy policy. A.D.A.M. is also a founding member of Hi-Ethics and subscribes to the principles of the Health on the Net Foundation (www.hon.ch).

The information provided herein should not be used during any medical emergency or for the diagnosis or treatment of any medical condition. A licensed medical professional should be consulted for diagnosis and treatment of any and all medical conditions. Call 911 for all medical emergencies. Links to other sites are provided for information only -- they do not constitute endorsements of those other sites. © 1997- 2011 A.D.A.M., Inc. Any duplication or distribution of the information contained herein is strictly prohibited.

