


[AUA Member Login](#) | [Login](#) | [Register](#) | [Claim Subscription](#) | [Subscribe](#)
[Articles & Issues](#)[Collections](#)[News & Media](#)[For Authors](#)[Journal Info](#)[AUA](#)[Related Sites](#)[Submit Your Manuscript](#)[Search AUA Journals](#)[All Content](#)[Search](#) [Advanced Search](#)[Share](#)< Previous ArticleMay 2016 Volume 195, Issue 5, Pages 1550–1555Next Article >Access this article on
[ScienceDirect](#)

To read this article in full, please review your options for gaining access at the bottom of the page.

Visit ScienceDirect to see if
you have access via your
institution.

Penile Low Intensity Shock Wave Treatment is Able to Shift PDE5i Nonresponders to Responders: A Double-Blind, Sham Controlled Study

Noam D. Kitrey*, , Ilan Gruenwald*, Boaz Appel†, Arik Shechter*, Omar Massarwa*, Yoram Vardi‡

Re: Therapeutic Hypothermia in Deceased Organ Donors and Kidney-Graft Function Goldfarb, David A.

[j.juro.2015.12.049](#)

[Abstract](#) [Full Text](#) [Images](#) [References](#)

Purpose

We performed sham controlled evaluation of penile low intensity shock wave treatment effect in patients unable to achieve sexual intercourse using PDE5i (phosphodiesterase type 5 inhibitor).

Materials and Methods

This prospective, randomized, double-blind, sham controlled study was done in patients with vasculogenic erectile dysfunction who stopped using PDE5i due to no efficacy. All patients had an erection hardness score of 2 or less with PDE5i. A total of 58 patients were randomized, including 37 treated with low intensity shock waves (12 sessions of 1,500 pulses of 0.09 mJ/mm² at 120 shock waves per minute) and 18 treated with a sham probe. In the sham group 16 patients underwent low intensity shock wave treatment 1 month after sham treatment. All patients were evaluated at baseline and 1 month after the end of treatment using validated erectile dysfunction questionnaires and the flow mediated dilatation technique for penile endothelial function. Erectile function was evaluated while patients were receiving PDE5i.

Results

In the low intensity shock wave treatment group and the sham group 54.1% and 0% of patients, respectively, achieved erection hard enough for vaginal penetration, that is an EHS (Erection Hardness Score) of 3 ($p < 0.0001$). According to changes in the IIEF-EF (International Index of Erectile Function-Erectile Function) score treatment was effective in 40.5% of men who received low intensity shock wave treatment but in none in the sham group ($p = 0.001$). Of patients treated with shock waves after sham treatment 56.3% achieved erection hard enough for penetration ($p < 0.005$).

Conclusions

Low intensity shock wave treatment is effective even in patients with severe erectile dysfunction who are PDE5i nonresponders. After treatment about half of them were able to achieve erection hard enough for penetration with PDE5i. Longer followup is needed to establish the place of low intensity shock wave treatment in these challenging cases.

[Create Citation Alert](#) [Cited by in Scopus](#) [Request Permissions](#) [Order Reprints](#) (100 minimum order)

Current Practices of Measuring and Reference Range Reporting of Free and Total Testosterone in the United States Le, Margaret et al.

Related Articles

[Low Intensity Shock Wave Treatment for Erectile Dysfunction—How Long Does the Effect Last?](#)

Noam D. Kitrey, Yoram Vardi, Boaz Appel, Arik Shechter, Omar Massarwi, Yasmin Abu-Ghanem, and others

[Shock Wave Treatment for Stones in the Upper Urinary Tract](#)

C. Chaussy, E. Schmiedt

[High Energy Underwater Shock Wave Treatment on Implanted Urinary Bladder Cancer in Rabbits](#)

Senji Hoshi, Seiichi Orikasa, Masa-Aki Kuwahara, Kenichi Suzuki, Kazuyuki

Key Words:

[testis](#), [erectile dysfunction](#), [high energy shock waves](#), [phosphodiesterase 5 inhibitors](#), [questionnaires](#)

Abbreviation and Acronyms:

[CGIC \(Clinical Global Impression of Change\)](#), [ED \(erectile dysfunction\)](#), [EF \(erectile function\)](#), [FMD \(flow mediated dilatation\)](#), [LIST \(low intensity shock wave treatment\)](#), [MCID \(minimal clinically important difference\)](#)

To access this article, please choose from the options below

Log In

[Login to existing account](#)

[Forgot password?](#)

Register

[Create a new account](#)

Purchase access to this article

- [\\$35.95 USD|PDF Download and 24 Hours Online Access](#)

Claim Access

If you are a current subscriber with Society Membership or an Account Number, [claim your access now.](#)

Subscribe to this title

[Purchase a subscription](#) to gain access to this and all other articles in this journal.

Institutional Access

[Visit ScienceDirect](#) to see if you have access via your institution.

Yoshikawa, Seiichi Saitoh, and others

Injury of Rat Renal Vessels Following Extracorporeal Shock Wave Treatment

Christian Weber, Michael E. Moran, Eldon J. Braun, George W. Drach

Use of Extracorporeal Shock Waves in the Treatment of Pseudarthrosis, Tendinopathy and Other Orthopedic Diseases

Gerald Haupt

[View All](#)

The corresponding author certifies that, when applicable, a statement(s) has been included in the manuscript documenting institutional review board, ethics committee or ethical review board study approval; principles of Helsinki
Re: Therapeutic Hypothermia in Deceased Organ Donors and Kidney-Graft Function Goldfarb, David A.

lieu of formal ethics committee approval; institutional animal care and use committee provided written informed consent with guarantees of confidentiality; IRB approved project number.

restricted grant from Medispec Ltd., Israel.

Current Practices of Measuring and Reference Range Reporting of Free and Total Testosterone in the United States Le, Margaret et al.

* No direct or indirect commercial incentive associated with publishing this article.

† Financial interest and/or other relationship with Medispec.

© 2016 American Urological Association Education and Research, Inc. Published by Elsevier Inc. All rights reserved.

[< Previous Article](#)

May 2016 Volume 195, Issue 5, Pages 1550–1555

[Next Article >](#)

Copyright © 2018 Elsevier Inc. All rights reserved. | [Privacy Policy](#) | [Terms & Conditions](#) | [Use of Cookies](#) | [About Us](#) | [Help & Contact](#) | [Accessibility](#)

The content on this site is intended for health professionals.

Advertisements on this site do not constitute a guarantee or endorsement by the journal, Association, or publisher of the quality or value of such product or of the claims made for it by its manufacturer.