



The first and only
FDA-approved
treatment for Thyroid
Eye Disease (TED)



TEPEZZATM
teprotumumab-trbw

Learn more at TEPEZZA.com

TEPEZZA is a prescription medicine used to
treat Thyroid Eye Disease.

**Please see Important Safety Information on
pages 28-31 and Full Prescribing Information
at TEPEZZA.com.**

Thyroid Eye Disease has many symptoms and can affect your life in different ways

Thyroid Eye Disease is a serious, vision-threatening condition that can get worse over time. If it is not treated early enough, it can lead to permanent damage to the eyes.

So, the sooner you and your doctor make a treatment plan, the better.

In this brochure, you will learn more about the disease, and how TEPEZZA (teprotumumab-trbw) may help.

Table of Contents

Facts about Thyroid Eye Disease.....	4
How TEPEZZA works.....	8
How TEPEZZA relieves symptoms.....	10
TEPEZZA and daily life.....	14
How TEPEZZA is given.....	16
Connect with a Nurse Advocate.....	18
Talking to your doctor	20
Important Safety Information.....	28

Select Important Safety Information

The most common side effects of TEPEZZA include muscle cramps or spasms, nausea, hair loss, diarrhea, feeling tired, high blood sugar, hearing problems, taste changes, headache, and dry skin. 9 out of 10 people who were treated with TEPEZZA in clinical studies completed all 8 doses.

Please see Important Safety Information on pages 28-31 and Full Prescribing Information at [TEPEZZA.com](https://www.tepezza.com).

Thyroid Eye Disease and Graves' disease are different and need different treatments

Up to 50% of people who have Graves' disease will develop Thyroid Eye Disease (TED). Despite this connection, Graves' disease and TED are separate conditions that affect different parts of the body.

Graves' disease affects your thyroid and TED affects your eyes.

That's why each condition needs to be treated differently.

Read on to learn more about the specialists who treat TED. Find one at TEDspecialist.com

The FDA classified TEPEZZA as a breakthrough medicine that treats TED


TEPEZZA[™]
teprotumumab-trbw

Select Important Safety Information

If you have inflammatory bowel disease (IBD), such as Crohn's disease or ulcerative colitis, TEPEZZA may make your IBD symptoms worse. Symptoms of worsening IBD may include: an increased number of loose stools with stomach pain or cramps, and blood in your stools. After each TEPEZZA infusion, tell your doctor right away if you have worsening IBD symptoms.

Act now to avoid permanent damage


Thyroid Eye Disease (TED) is a progressive disease, which means it gets worse over time. When you first develop TED, it is progressing (or “active”). During this time, symptoms like eye bulging, pain, redness, and swelling appear, and can change or continue to get worse. The progressing (or “active”) phase of TED can last for as little as 6 months or up to 3 years.

Over time, the progressing (or “active”) phase of TED changes, and scar tissue begins to form. During this non-progressing (or “inactive”) phase, symptoms such as redness and swelling may get better. Even though symptoms may improve, there could be permanent damage to the eyes.

Get treatment for TED as early as possible, before damage becomes permanent

Please see Important Safety Information on pages 28-31 and Full Prescribing Information at [TEPEZZA.com](https://www.tepezza.com).

What happens during the progressing (or “active”) phase of TED:



You may also notice:




TEPEZZA is the only medicine that treats the cause, not just the symptoms, of Thyroid Eye Disease

Unlike other medicines you and your doctor may have tried, TEPEZZA (teprotumumab-trbw) treats Thyroid Eye Disease (TED) at the source to reduce bulging eyes, double vision, eye pain, redness, and swelling.

Select Important Safety Information


TEPEZZA may cause an increase in your blood sugar. Before starting treatment with TEPEZZA, tell your doctor if you are currently being treated for diabetes, know your blood sugar is high, or have been diagnosed with diabetes. It is important for you to take your treatments and follow an appropriate diet for glucose control as prescribed by your doctor.

How TED affects the eyes



A special switch called a receptor gets turned on. This leads to swelling of the muscle and fat tissue behind your eyes, causing symptoms like eye bulging, double vision, pain, and redness.

How TEPEZZA works



TEPEZZA blocks this receptor. By blocking the receptor, TEPEZZA reduces muscle and fat tissue swelling, and improves symptoms such as bulging eyes, double vision, eye pain, redness, and swelling.

Please see Important Safety Information on pages 28-31 and Full Prescribing Information at [TEPEZZA.com](https://www.tepezza.com).

Look more like your old self

In a clinical study, more than

8 out of 10

people taking TEPEZZA (teprotumumab-trbw) saw reduced eye bulging.*

In another study, **more than half** of the people who had less eye bulging* after completing treatment with TEPEZZA maintained their improvements for close to a year.



With TEPEZZA, some people began to see less eye bulging in as few as 6 weeks, with improvements continuing for the full course of treatment. And no one took steroids during their treatment.

*Eye bulging was reduced by at least 2 mm.

Select Important Safety Information

The most common side effects of TEPEZZA include muscle cramps or spasms, nausea, hair loss, diarrhea, feeling tired, high blood sugar, hearing problems, taste changes, headache, and dry skin. 9 out of 10 people who were treated with TEPEZZA in clinical studies completed all 8 doses.

Before and after treatment with TEPEZZA



Real patient treated with TEPEZZA for 24 weeks in a clinical study. Individual results may vary.

Please see Important Safety Information on pages 28-31 and Full Prescribing Information at [TEPEZZA.com](https://www.tepezza.com).

See more clearly



In clinical studies,

7 out of 10

people taking TEPEZZA (teprotumumab-trbw) saw improvements in double vision.



In clinical studies, more than

5 out of 10

people taking TEPEZZA saw their double vision completely go away.

In another study, **7 out of 10** people who had improvements in double vision still had no double vision almost a year after finishing treatment.

Please see Important Safety Information on pages 28-31 and Full Prescribing Information at [TEPEZZA.com](https://www.tepezza.com).

Get multi-symptom relief

In a clinical study,

2 out of 3

people taking TEPEZZA had little to no:



Eye Pain

including behind the eye and when moving the eye



Redness

of eye/eyelid



Swelling

of eye/eyelid


Some people had less eye bulging, pain, redness, and swelling starting as early as 6 weeks, and improvement continued throughout treatment with TEPEZZA.

Select Important Safety Information

Infusion reactions can happen during or within 24 hours after your infusion of TEPEZZA. If you have a reaction while receiving TEPEZZA, your doctor or nurse will slow or stop your infusion and treat your reaction. If you have a severe infusion reaction, your doctor may stop your treatment completely.

In clinical studies of people
with Thyroid Eye Disease,

TEPEZZA improved visual ability



**Please see Important Safety Information on
pages 28-31 and Full Prescribing Information
at [TEPEZZA.com](https://www.tepezza.com).**

And physical appearance



Select Important Safety Information

Before receiving TEPEZZA (teprotumumab-trbw), tell your doctor if you are pregnant or plan to become pregnant. TEPEZZA may harm your unborn baby. Tell your doctor if you become pregnant or suspect you are pregnant during treatment with TEPEZZA. Women who are able to become pregnant should use an effective form of birth control (contraception) prior to starting treatment, during treatment and for at least 6 months after the final dose of TEPEZZA.

How TEPEZZA is given

TEPEZZA (teprotumumab-trbw) is an IV (intravenous) medicine. This means it is delivered through a needle that is placed in your arm.


IV treatments like TEPEZZA are most often given by registered nurses (RNs), nurse practitioners (NPs), or another medical professional.

The amount of TEPEZZA that you will receive is based on how much you weigh.

Select Important Safety Information

Infusion reactions can happen during or within 24 hours after your infusion of TEPEZZA. If you have a reaction while receiving TEPEZZA, your doctor or nurse will slow or stop your infusion and treat your reaction. If you have a severe infusion reaction, your doctor may stop your treatment completely.

With just 8 doses—1 dose every 3 weeks for about 5 months—you may experience fewer symptoms of Thyroid Eye Disease



Your first and second treatments of TEPEZZA will last about 90 minutes each. Your remaining 6 treatments will take about 60 minutes each, but could be longer depending on how you react to the infusion.

To get the full benefit of TEPEZZA, it is important that you complete all 8 doses

Please see Important Safety Information on pages 28-31 and Full Prescribing Information at [TEPEZZA.com](https://www.tepezza.com).

Talk to a Nurse Advocate

Whether you are newly diagnosed or already have experience with Thyroid Eye Disease (TED), Nurse Advocates are here to help you:

- Learn more about TED
- Discuss your treatment options, including TEPEZZA (teprotumumab-trbw)
- Find a TED specialist near you



Call 1-844-4-TEPEZZA
Monday-Friday 8 AM-8 PM, ET

Nurse Advocates do not provide individual patient care or medical advice. You should talk to your doctor about whether TEPEZZA is right for you.

Please see Important Safety Information on pages 28-31 and Full Prescribing Information at [TEPEZZA.com](https://www.tepezza.com).

Meet the TED specialists

TED is complicated and rare, and not all doctors are used to treating it. That is why you should see a team of doctors who specialize in treating TED. This team should be made up of:

Eye specialists (ophthalmologist/oculoplastic surgeon) will:

- Give you an eye exam
- Determine if your TED is in the progressing (or “active”) or non-progressing (or “inactive”) phase
- Create a treatment plan
- Work with your endocrinologist to make sure your TED treatment does not affect any medicines you may be taking for another thyroid condition, like Graves’ disease

Endocrinologists will:

- Manage any thyroid condition you may have, like Graves’ disease
- Work with your eye specialist on a treatment plan for TED

To find TED specialists in your area, visit TEDspecialist.com

Be honest about how Thyroid Eye Disease impacts your life

The more your doctors know about how Thyroid Eye Disease (TED) affects you and your daily life, the more they can help.

TED doesn't just affect your eye health. The physical and emotional changes you may experience are just as important—and should not be ignored.

Before your doctor appointments, you may find it helpful to:

- 1.** Track the changes to your eyes
- 2.** Keep a record of your day-to-day life
- 3.** Write down questions you want to ask your doctor

With this information, you and your doctor can work together to develop the best treatment plan for you.

**Use the following 3-step guide
to get started**



Step 1: Track your symptoms

Over the past month, how often have you experienced the following Thyroid Eye Disease (TED) symptoms? (Check the box that applies.)

	ALL THE TIME	SOMETIMES	NEVER
Eye bulging	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Double vision	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Eye pain	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Redness in/around the eye	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Swelling in/around the eye	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Use the space below to write in any other TED-related symptoms that may be bothering you.

Tips to keep track of your symptoms



Take selfies
Taking selfies is a great way to keep track of how your symptoms and appearance are changing over time. Try to zoom in on your eyes when you take your selfie.



Keep a symptom journal
Recording symptoms in a journal is another helpful way to track changes. Be specific about the types of TED symptoms you notice and how they impact your life.



Create a routine
It is important to check your eyes often. Having a routine can help you remember. So, try picking a day of the week to check your eyes.

If you see any changes to your eyes, speak to a TED specialist right away

Please see Important Safety Information on pages 28-31 and Full Prescribing Information at TEPEZZA.com.

Step 2: Keep a record of your day-to-day life

Daily life

Over the past month, has Thyroid Eye Disease (TED) made it harder to do any of these daily tasks? (Circle the answer that applies.)

Walking outdoors	Yes	No	Not sure
Reading	Yes	No	Not sure
Driving	Yes	No	Not sure
Watching TV	Yes	No	Not sure
Cooking	Yes	No	Not sure
Working	Yes	No	Not sure
Walking up or down stairs	Yes	No	Not sure

Are there any other activities or hobbies that you wanted to do but could not because of TED? (Use the space below to write in your answer.)

Wellbeing

Over the past month, have you noticed any of the following? (Circle the answer that applies.)

Depressive feelings	Yes	No	Not sure
Anxiety	Yes	No	Not sure
Feeling like you are being watched or judged	Yes	No	Not sure
Hiding your appearance in public	Yes	No	Not sure

Are there other ways that TED has changed how you feel about yourself? (Use the space below to write in your answer.)

Step 3: Talk to your doctor about TEPEZZA

Here are some questions your doctor can answer for you, so you can both decide if TEPEZZA (teprotumumab-trbw) is the right treatment for you:

- How is TEPEZZA different from other treatments I have tried?
- How could TEPEZZA reduce my symptoms?
- What are the possible side effects of TEPEZZA?
- How will I receive TEPEZZA?
- How long will I receive treatment?
- Where will I receive treatment?

Select Important Safety Information

Infusion reactions can happen during or within 24 hours after your infusion of TEPEZZA. If you have a reaction while receiving TEPEZZA, your doctor or nurse will slow or stop your infusion and treat your reaction. If you have a severe infusion reaction, your doctor may stop your treatment completely.

Use the space below to write in any other questions you may have.

Please see Important Safety Information on pages 28-31 and Full Prescribing Information at [TEPEZZA.com](https://www.tepezza.com).

USE

TEPEZZA is a prescription medicine used to treat Thyroid Eye Disease.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about TEPEZZA?

Infusion reactions can happen during or within 24 hours after your infusion of TEPEZZA. If you have a reaction while receiving TEPEZZA, your doctor or nurse will slow or stop your infusion and treat your reaction. If you have a severe infusion reaction, your doctor may stop your treatment completely.

Tell your doctor or nurse right away if you have any of these symptoms during or after your treatment with TEPEZZA:

- High blood pressure
- Fast heartbeat
- Redness of the face/Feeling hot
- Difficulty breathing
- Headache
- Muscle pain

If you have inflammatory bowel disease (IBD), such as Crohn's disease or ulcerative colitis, TEPEZZA may make your IBD symptoms worse. Symptoms of worsening IBD may include: an increased number of

loose stools with stomach pain or cramps, and blood in your stools. After each TEPEZZA infusion, tell your doctor right away if you have worsening IBD symptoms.

TEPEZZA may cause an increase in your blood sugar. Before starting treatment with TEPEZZA, tell your doctor if you are currently being treated for diabetes, know your blood sugar is high, or have been diagnosed with diabetes. It is important for you to take your treatments and follow an appropriate diet for glucose control as prescribed by your doctor.

Before receiving TEPEZZA, tell your doctor if you:

- Have inflammatory bowel disease (Crohn's disease or ulcerative colitis).
- Are currently being treated for diabetes, have been diagnosed with diabetes, or know your blood sugar is high.
- Are pregnant or plan to become pregnant. TEPEZZA may harm your unborn baby. Tell your doctor if you become pregnant or suspect you are pregnant during treatment with TEPEZZA.

Please see Full Prescribing Information at [TEPEZZA.com](https://www.tepezza.com).

IMPORTANT SAFETY INFORMATION (continued)

- Women who are able to become pregnant should use an effective form of birth control (contraception) prior to starting treatment, during treatment and for at least 6 months after the final dose of TEPEZZA.
- Are breastfeeding or plan to breastfeed. It is not known if TEPEZZA passes into your breast milk. Talk to your doctor about the best ways to feed your baby during treatment with TEPEZZA.
- Tell your doctor about all the medicines you take, including prescription and over the counter medicines, vitamins, dietary and herbal supplements. Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

What are the possible side effects of TEPEZZA?

The most common side effects of TEPEZZA include muscle cramps or spasms, nausea, hair loss, diarrhea, feeling tired, high blood sugar, hearing problems, taste changes, headache, and dry skin.

This is not a complete list of all possible side effects. Tell your doctor or treatment team if you have any side effect that bothers you or that does not go away.

Please visit [TEPEZZA.com](https://www.tepezza.com) for more information.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/safety/medwatch, or call the FDA at [1-800-FDA-1088](tel:1800FDA1088).

Please see Full Prescribing Information at [TEPEZZA.com](https://www.tepezza.com).



See the difference TEPEZZA can make

TEPEZZA (teprotumumab-trbw) provides relief from **eye bulging, double vision, redness, swelling, and pain.**

For more information, visit TEPEZZA.com and talk to your TED specialist

Select Important Safety Information

The most common side effects of TEPEZZA include muscle cramps or spasms, nausea, hair loss, diarrhea, feeling tired, high blood sugar, hearing problems, taste changes, headache, and dry skin. 9 out of 10 people who were treated with TEPEZZA in clinical studies completed all 8 doses.

Please see Important Safety Information on pages 28-31 and Full Prescribing Information at TEPEZZA.com.