

USE OF PLANT BASED STEM CELL NANOTECHNOLOGY ON NONHEALING CHRONIC WOUNDS

Naz Wahab MD, FAAP, FAPWCA¹, Jutta A. Guadagnoli PhD^{2,3}, Kelli Wray APRN, AGNP-C¹, and Tammy

Luttrell MSPT, PhD, CWS, FCCWC⁴

¹Wound Care Associates, Las Vegas, NV; ²Department of Clinical Education, Touro University Nevada School of Osteopathic Medicine, Henderson, NV; ³Department of Basic Sciences, Touro University Nevada School of Osteopathic Medicine, Henderson, NV; ⁴Department of Physical Therapy University Medical Center Nevada

INTRODUCTION

Advanced wound care products for chronically non-healing wounds have been a growing area of wound healing technology. As the science behind wound healing becomes more widely investigated newer stem cell therapies are being brought onto the market. The numbers of non-healing diabetic wounds are growing due to complex multi-organism bio-burden, advancing diabetic co-morbidities, and limitations of cost effective advanced therapies. Stem cell therapy has exploded on the forefront of wound healing products, but most are of human, or other animal origin. Isolating stem cells from animal origins is a complex, and costly approach which leads to difficulty in delivering advanced modalities to patients in need.

An alternative was to isolate undifferentiated plant stem cells from green tea and create a hemicellulose membrane with Nano particulate fibers. This membrane using 100% non-animal ingredients with the small nanotechnology delivery system is a successful approach to delivering stem cells directly to the wound bed.


It is known that products that have animal undifferentiated stem cells can be effective in wound healing, but are plant stem cells effective as well? If the chronically stalled diabetic wound could respond by effectively decreasing wound size and volume then it could be a viable treatment option.

METHODS




The green tea membrane was applied onto 32 patients with chronic non-healing wounds of various etiologies that did not improve greater than 30% with decrease in wound size or volume for a minimum of 6 weeks. These patients were identified and serially categorized from the study initiation date. Secondary dressings of untreated gauze or foam were utilized. Wound size and closure rates were monitored over time. Adverse reactions associated with the treatment regime were noted.

RESULTS

- Total number of wounds 32
- Mean age 62.5 ± 14.5 years
- Three did not complete the study due to loss of follow up, infection, or Physician discretion
- Avg. duration of care 7 weeks
- Avg. heal rate for patients who healed completely was 20.3% reduction/wk
- Avg. number of weeks to heal with nanogen 4.8 weeks
- Avg. wound age prior to starting therapy was 10.8months +1.5 months
- 59% diabetic wounds; 32% pressure related wounds, other etiologies radiation induced, surgical, and autoimmune wounds



Heal Rate: At 7 weeks 75% of patients healed although not all healed at the end of the study



DISCUSSION

The Plant Stem Cell Nanotechnology is a 100% natural biological hemicellulose membrane. It is a green tea based fiber scaffold of nanometric sized particles creating a plant stem cell nanodevice. This nanodevice is an innovative way of delivering stem cells directly into the wound bed. Stem cell nanoparticles have the ability to enter into the cytoplasmic space across cellular barriers and activate specific transport mechanisms with direct cell to cell signaling based on the combination of both quantum particle size and the on the presence of the undifferentiated plant stem cells. This nanodevice membrane helps to replace the extracellular matrix by regulating growth factors, direct cell signaling, and cell to matrix signaling.

When the green tea nanodevice membrane is applied to the wound bed, deposition of extracellular matrix occurs due to the direct chemotactic signaling of platelets, macrophages, T lymphocytes, and fibroblasts. These cell types in turn produce PDGF, TGF- β , EGF, FGF, and IGF. The initiation of these growth factors then stimulates the proliferative phase of wound healing. More specifically, fibroblasts will release collagen and glycosaminoglycans, which in combination with fibronectin, forms the new extracellular matrix. The body's natural ECM then is the framework for angiogenesis, and continued granulation tissue formation and finally keratinocyte migration and epithelialization.

The green tea nanodevice membrane also carries a positive charge due to the amine groups on the highly pedicted carboxylic acid cellulose. The positive charge creates an electromagnetic attraction with the negatively charged blood byproducts and negatively charged bacteria. Therefore, upon wound bed stimulation by the membrane it is clinically and qualitatively seen that deep-seated bio burden is removed and edema is improved.

The benefits of this wound healing modality are far reaching. The cell signaling is faster and more effective due to the nanotechnology delivery system. The medicinal value of the plant itself can be utilized directly for wound healing. The products are 100% bio-sustainable and biodegradable.

The expense in collection, isolation, and purification of the animal based products can be timely and expensive. Utilizing plants instead of animals can be a more cost effective alternative. Plant based products can also be less immunogenic due to lack of species cross reactivity. There is no clinical preparation, no difficult storage limitations, greater shelf life, greater patient selection versatility, and overall more cost effective.

REFERENCES

1. G. A. Olywinski, D. D. Vukelic, L. M. W. Costa, R. N. Seetharam, L. X. Filipe, P. Benavelli, First-Order Cellular/Nanobiological Cellulose Nanocomposites as a Potential Scaffold for Bone Tissue Regeneration, *J. Biomater. Nanotechnol.* 2, 239-251 (2011).
2. Novel antimicrobial peptides bacterial cellulose obtained by biomass culture between poly(hydroxybutyrate) (PHB) and green tea *Pithecellobium dulce* (PDC) leaves, *Journal of Biomedical Materials Research Part B: Applied Biomaterials*, 93B, 100-108 (2011).
3. Pisanelli, G.M., Olywinski, L.M.W. Costa, L.X. Filipe, Bacterial Nanocellulose for medicine regenerative, *J. Microtech. Eng. Med.* 2, 024001, 2011.
4. R. N. Seetharam, "Nanomedicine—emerging area of nanotechnology research," *Current Science*, vol. 91, no. 3, p. 260, 2011.