

Medication hold list for surgery

- | | | | |
|---|---------------------|------------------|----------------------|
| • Aggrenox | • Combunox | • Ketoprofen | • PMS-ASA |
| • Aleve | • Cope | • Ketorolac | • Ponstel |
| • Alka-Seltzer | • Daypro | • Lodine | • Prevacid NapraPAC |
| • Amigesic | • Diclofenac | • Lovaza | • Relafen |
| • Anacin products | • Diflunisal | • Magan | • Robaxisal |
| • Anaflex | • Disalcid | • Magnaprin | • Roxiprin |
| • Anaprox | • Doan's | • Marthritic | • Salflex |
| • Ansaid | • Dolobid | • Meclofenamate | • Salsalate |
| • Apo-ASEN | • Easprin | • Meclomen | • Sine-Aid IB |
| • Arco Pain Tablet | • Ecotrin 81 | • Medipren | • Sodium salicylate |
| • Argeric | • Empirin | • Mefenamic acid | • Soma Compound |
| • Arthropan | • Endodan | • Midol | • St. Joseph Aspirin |
| • Arthrotec | • Entrophen | • Mobic | • Sulindac |
| • Ascriptin | • Equagesic | • Mobiflex | • Suprofen |
| • Aspergum | • Es Anacin | • Momentum | • Suprol |
| • Aspirin (all products containing aspirin) | • Etodolac | • Mono-Gesic | • Surgam |
| Aspir-Low | • Excedrin Migraine | • Motrin | • Synalgos-DC |
| • Aspartab | • Feldene | • Nabumetone | • Tandearil |
| • Bayer | • Fenoprofen | • Nalfon | • Talwin Compound |
| • Bayer time release | • Fiorinal | • Naprelan | • Tenoxicam |
| • Buffex | • Flector Patch | • Naprosyn | • Tiaprofenic acid |
| • Bufferin | • Floctafenine | • Naproxen | • Tolectin |
| • Bufferin Arthritis | • Flurbiprofen | • Norgesic Forte | • Tolmetin |
| • Buffinol | • Glucosamine | • Nuprin | • Toradol |
| • Butalbital Compound | • Goody's | • Ocuville | • Tricosal |
| • Butazolidin | • Halfprin | • Oruvail | • Trilisate |
| • Cama Arthritis | • Helidac | • Orudis | • Vanquish |
| • Carisoprodol Compound | • Ibuprofen | • Oxaprozin | • Vicoprofen |
| • Cataflam | • Indocin | • Pamperin-IB | • Voltaren |
| • Clinoril | • Indomethacin | • Pepto-Bismol | • Zorprin |
| | • Instantine | • Percodan | |
| | • Kaopectate | • Phenylbutazone | |
| | • Kava | • Piroxicam | |

We recommend stopping **Aggrenox, Arixtra, Aspirin, Brilinta, Coumadin, Eliquis, Fragmin, Innohep, Lovenox, Plavis, Pletal, Pradax** and **Xarelto** with approval of your prescribing physician to have normal coagulation at the time of surgery. We also recommend stopping supplements and other over the counter agents 7 days prior to surgery.

By signing, I agree that I must not take any of these over-the-counter medications for the time frame specified. I understand that failure to follow these instructions might result in the postponement of my surgery.

Name of Patient/Legal Representative

Signature

Date