



OrthoTeam Clinic

Ashish M. Rawal, M.D.

2 Science Court, Madison, WI 53711

900 Ridge Street, Stoughton, WI 53589.

Telephone: 608-231-3410 Fax: 608-231-3430

Post-Op Instructions ACL Reconstruction with Meniscal Repair

DIET

- Begin with clear liquids and light foods (Jell-O, soups, etc.)
- Progress to your normal diet if you are not nauseated

WOUND CARE

- Maintain your operative dressing, and loosen bandage if swelling of the foot and ankle occurs
- It is normal for the knee to bleed and swell following surgery. If blood soaks onto the TED stocking bandage, do not become alarmed, reinforce with additional dressings as needed.
- To avoid infection, keep surgical incisions clean and dry for the first **7 days** following surgery – you may shower by placing a large plastic bag over your brace beginning the day after surgery. NO immersion of the operative leg (ie: bath or pool).
- Wait until your first post-operative appointment to have Dr. Rawal's team remove the surgical dressing
- Please do not place any ointments, lotions, or creams directly over the incisions.
- Once the sutures are removed **at least 7-14 days post operatively** you can begin to get the incision wet in the shower (water and soap lightly run over the incision and pat dry). NO immersion in a bath until given approval by our office.

MEDICATIONS

- Local anesthetics are injected into the wound and knee joint at the time of surgery. This will wear off within 12-24 hours and it is not uncommon for patients to encounter more pain on the first or second day after surgery when swelling peaks.
- Most patients will require some narcotic pain medication for a short period of time – this can be taken as per directions on the bottle.
- Common side effects of the pain medication are nausea, drowsiness, and constipation. To decrease these side effects, take the medication with food. If constipation occurs, consider taking an over the counter laxative and be sure to drink plenty of water.
- If you are having problems with nausea and vomiting, contact the office to possibly have your medications changed.
- Do not drive a car or operate machinery while taking the narcotic medication.
- Please avoid alcohol use while taking narcotic pain medication.
- If you are having pain that is not being controlled by the pain medication prescribed, you may take an over the counter acetaminophen (Tylenol) as directed on the bottle. This will help to decrease pain and decrease the amount of narcotic medication required.
- For 4 weeks following surgery take **one aspirin 325mg tablet** daily to lower the risk of developing a blood clot after surgery. Please contact the office should severe calf pain occur or significant swelling of the calf or ankle occur.



OrthoTeam Clinic

Ashish M. Rawal, M.D.

2 Science Court, Madison, WI 53711

900 Ridge Street, Stoughton, WI 53589.

Telephone: 608-231-3410 Fax: 608-231-3430

ACTIVITY

- Elevate the operative leg to chest level whenever possible to decrease swelling.
- Do not place pillows under knees (i.e. do not maintain knee in a flexed or bent position), but rather place pillows under the foot/ankle.
- Due to the nature of the meniscal repair, you will be **Touch Down Weight Bearing** for balance (ie non weight bearing with crutches but only resting the foot on the ground for balance) **for 4-6 weeks** post operatively
- Do not engage in activities which increase knee pain/swelling (prolonged periods of standing or walking) for the first 7-10 days following surgery.
- Avoid long periods of sitting (without leg elevated) or long distance traveling for 2 weeks.
- NO driving until instructed otherwise by physician
- May return to sedentary work ONLY or school 3-4 days after surgery, if pain is tolerable

BRACE

- Your brace should be worn fully extended (straight) at all times (day and night – except for exercises) until otherwise instructed after the first post-operative visit.

ICE THERAPY

- Icing is very important in the initial post-operative period and should begin immediately after surgery.
- Use icing machine continuously or ice packs (if machine not prescribed) for 45 minutes every 2 hours daily until your first post-operative visit. Care should be taken with icing to avoid frostbite to the skin. **DO NOT PLACE ICE PACK / PAD DIRECTLY ON SKIN.**
- Icing should not be performed while sleeping.

EXERCISE

- Begin exercises 24 hours after surgery (straight leg raises, quad sets, heel slides, and ankle pumps) unless otherwise instructed.
- Discomfort and knee stiffness is normal for a several days following surgery. It is safe to bend your knee in a non-weightbearing position when performing exercises unless otherwise instructed. Avoid flexing past 90 degrees. Bending beyond 90 degrees is not recommended until 4-6 weeks after surgery.
- Complete exercises 3-4 times daily until your first post-operative visit – your motion goals are to have complete extension (straightening) and 90 degrees of flexion (bending) at your first post-operative appointment unless otherwise instructed.
- Perform ankle pumps continuously throughout the day to reduce the risk of developing a blood clot in your calf.
- Formal physical therapy (PT) typically begins a few days after surgery. A PT rehab protocol will be provided at your pre-operative visit.



OrthoTeam Clinic

Ashish M. Rawal, M.D.

2 Science Court, Madison, WI 53711

900 Ridge Street, Stoughton, WI 53589.

Telephone: 608-231-3410 Fax: 608-231-3430

EMERGENCIES**

- Contact our office at (608) 231-3410 if any of the following are present:
- Painful swelling or numbness (note that some swelling and numbness is normal)
- Unrelenting pain
- Fever (over 101° – it is normal to have a low grade fever for the first day or two following surgery) or chills
- Redness around incisions
- Color change in foot or ankle
- Continuous drainage or bleeding from incision (a small amount of drainage is expected)
- Difficulty breathing
- Excessive nausea/vomiting
- Calf pain
- If you have an emergency that requires immediate attention proceed to the nearest emergency room.

FOLLOW-UP CARE/QUESTIONS

- If you do not already have a post-operative appointment scheduled, please contact our scheduler at 608-231-3410.
- Your first post-operative appointment will be scheduled with one of the Physician Assistants for a quick wound check, physical therapy protocol and to answer any further questions you have regarding the procedure
- Typically the first post-operative appointment following ACL reconstruction is made for 2-3 days following surgery and 10-14 days following surgery for suture removal.