


## Caudal Epidural Steroid Injection



### Overview

This outpatient procedure is an injection of a steroid-anesthetic medication through an opening in the sacrum. The medication can reduce swelling and inflammation of irritated spinal nerves. The injection takes only a few minutes to complete.

### Preparation

In preparation for the procedure, the patient lies face down. A cushion is placed under the abdomen to elevate the sacrum.

### Anesthetic Injected

The physician administers a local anesthetic to numb the skin and the tissue above the small opening at the base of the sacrum. This opening is called the sacral hiatus.

### Needle Inserted

When the area is numb, the physician guides a needle through the sacral hiatus and into the caudal epidural space. This is the open space in the sacrum where the irritated nerve roots are located.

### Contrast Solution Injected



The physician injects contrast solution through the needle. The physician uses a fluoroscope (a type of x-ray device) to confirm that the tip of the needle is positioned correctly within the epidural space.

### Medication Injected

After the needle's position has been confirmed, the physician injects a steroid-anesthetic medication. This medication bathes the irritated nerve roots. It will help alleviate the patient's pain.

### End of Procedure

When the procedure is complete, the physician removes the needle and bandages the insertion site. The patient may feel significant relief after one injection. Some patients may need multiple injections before they feel the full benefit of the medication.



1. If you are taking any blood thinners, you may continue to take these prior to your procedure. There is NO need to discontinue these medications.
2. Notify staff of any allergies: iodine, anesthetic, seafood, steroids, latex or contrast dye
3. Following your procedure, apply ice for 20 minutes every 1-2 hours.
4. If you encounter soreness after your procedure and are able to take NSAIDS (Ibuprofen, Aleve, Meloxicam) you may do so. Please follow medication directions on dosage and how often you may take these.

If you have any questions following your procedure or experience any complications please, call our office.