

Flexible Sigmoidoscopy & Pouchoscopy Instructions

Appropriate bowel cleansing is one of the **most important** steps in performing a successful and accurate sigmoidoscopy. Please read this sheet in advance of your procedure and follow these instructions carefully. If you have any questions please do not hesitate to call us.

One week prior to procedure:

- Purchase the following items at your local pharmacy:
 - One bottle of Magnesium Citrate - 10 ounce (296 mL)
 - Two Fleet Enemas
- If you take **blood thinners** such as Aspirin (or Aspirin containing products), Plavix (Clopidogrel), Effient (Prasugrel), Coumadin (Warfarin), Pradaxa (Dabigatran), Xarelto (Rivaroxaban), Eliquis (Apixaban), or any other blood thinners *please inform us in advance* so that these medicines can be adjusted appropriately.
- If you are a diabetic and take **insulin** or other medications to lower your blood sugar *please inform us in advance* so that these medicines can be adjusted appropriately.
- Stop any anti-inflammatory medicines (Aleve, Motrin, Advil, Ibuprofen, Naproxen). You can take Tylenol if needed.

One day before the procedure:

- You may have a regular breakfast and lunch. Starting in the early afternoon you should remain on a “clear/thin liquid” diet. Examples include water, broth, jell-o, juices, sports drinks, sodas, ginger ale, coffee/tea (*without* milk).
- Drink a bottle of Magnesium Citrate at 10 pm the evening before your procedure. This should produce bowel movements within a few hours. Drink plenty of fluids to maintain hydration and help with the cleansing process.

Day of the procedure:

- Approximately 2 to 4 hours before your procedure take a Fleets Enema and hold it for 10 minutes before evacuation.
- Repeat the process immediately using the second Fleet Enema.
- Arrive at your designated location at least **1 hour early** to allow for registration so that your procedure is not delayed.
- Do not drink anything within **4 hours** of your procedure time otherwise your procedure may be cancelled.
- You can take your medications (unless instructed otherwise) with water **4 hours** before your procedure. If you have an early morning appointment, you can bring your medications with you and take them after your procedure.
- You should be accompanied by a relative/friend to help you get home after the procedure.
- Women of child-bearing age may be asked to provide a urine sample to ensure you are not pregnant.
- You should not drive or drink alcohol for 12 hours after the procedure.

Procedure Locations:

	920 Park Avenue New York, NY 10028	T: (212) 535-3400
	Carnegie Hill Endoscopy Center 1516 Lexington Avenue (at East 98 th street) New York, NY 10029	T: (212) 860-6300
	Mount Sinai Hospital Endoscopy Unit 1184 5 th Avenue (at East 100 th street), 7 th floor New York, NY 10029	T: (212) 659-8961 (scheduling) T: (212) 241-6277 (endo unit)

Cancelling or Rescheduling:

- If you need to cancel or reschedule please call us at **212-535-3400** to inform us and allow other patients to use the procedure time.
- There is a \$50 fee for no-shows and late cancellations (less than 3 business days).

Please do not hesitate to call us with any questions!