

Hooman Khorasani, M.D. Chief

Division of Dermatologic and Cosmetic Surgery

**E85th Street Skin & Laser Center** 234 East 85<sup>th</sup> Street – 5<sup>th</sup> Floor New York, NY 10029-6574 Tel 212.731.3311 Fax 212.731.3395

## Vascular/Scar (VBeam) Laser Aftercare

- **1.** Immediately following the procedure you will notice a purple-colored bruise on the treated area. This will usually resolve after a period of up to 3 weeks.
- 2. Apply a cool compress to the treated area. This will help reduce any post-operative swelling and pain.
- a. Ice cubes in a bag, surrounded with a cool wash cloth if needed.
- **3.** Sleep with your head elevated on two pillows to reduce swelling around the eyes.
- 4. Once home, apply an emollient such as Aquaphor or Vaseline to keep the area moist.
- **5.** Tylenol is good for pain relief. Avoid Aspirins as they may break down any clots that are necessary to **ensure a good result.**
- **6.** Gentle, non-abrasive skin cleansing is recommended. DO NOT scrape the treated areas with items such as a rough wash cloth.
- 7. Should any blisters occur:
- a. You may gently pop them with a sterile pin or needle.
- b. DO NOT remove the blister roof. The extra skin is necessary to speed the healing of the skin.
- c. DO NOT pick at scabs or try to speed the healing process by removing the skin prematurely, this could result in scarring and pigment changes to the skin.
- 8. Sunscreens with at least SPF30 should be worn daily. Unprotected sun exposure should be avoided.
- 9. Re-treatments can be done at 6 to 8 week intervals.
- 10. Should there be any questions or concerns after the treatment please call your doctor.