



Fever Reducing Dosage Schedule

Children's Tylenol (Acetaminophen)

		Infant's Tylenol Oral Suspension	Children's Tylenol Oral Suspension	Children's Tylenol Soft Chews Chewable Tablets	Junior Strength Tylenol Chewable Tablets
Weight (lbs)	Age	160mg/5ml	160mg/5ml	80mg Tablet	160mg Tablet
6-11 lbs	2-3 months	1.25ml	_____	-----	-----
12-17 lbs	4-11 months	2.5ml	_____	-----	-----
18-23 lbs	12-23 months	3.75ml	_____	-----	-----
24-35 lbs	2-3 years	5ml	1 tsp=5ml	2	-----
36-47 lbs	4-5 years	-----	1 ½ tsp =7.5ml	3	-----
48-59 lbs	6-8 years	-----	2 tsp=10ml	4	2
60-71 lbs	9-10 years	-----	2 ½ tsp =12.5ml	5	2 ½
72-95 lbs	11 years	-----	3 tsp=15ml	6	3
>96 lbs	12 years	-----	-----	-----	4

- Never use Aspirin in children as it is associated with a dangerous condition called Reye's Syndrome
- Use only as directed by your physician
- You may give Tylenol every 4 to 6 hours but not more than 5 times in a 24 hour period
- Keep all medication out of the reach of children

Disclaimer: These guidelines are to help the caretaker with home treatment. If you are concerned about your child's health you should see your physician in person.



Children's Motrin (Ibuprofen)

		Infant's Motrin Concentrated Drops	Children's Motrin Suspension	Children's Motrin Chewable Tablets	Junior Strength Motrin Chewable Tablets	Junior Strength Motrin Caplets
Weight (lbs)	Age	50mg/1.25ml Dropperful	100mg/5ml Teaspoon	50mg Tablet	100mg Tablet	100mg Caplet
Under 12 lbs	Under 6 months	****DO NOT GIVE TO INFANTS UNDER 6 MONTHS OF AGE****				
12-17 lbs	6-11 months	1 dropper= 1.25ml	-----	-----	-----	-----
18-23 lbs	12-23 months	1 ½ dropper= 1.875ml	-----	-----	-----	-----
24-35 lbs	2-3 years	-----	1 tsp (5ml)	2	-----	-----
36-47 lbs	4-5 years	-----	1 ½ tsp (7.5ml)	3	-----	-----
48-59 lbs	6-8 years	-----	2 tsp (10ml)	4	2	2
60-71 lbs	9-10 years	-----	2 ½ tsp (12.5ml)	5	2 ½	2 ½
72-95 lbs	11 years	-----	3 tsp (15ml)	6	3	3
>96 lbs	12 years	-----	-----	-----	4	-----

- Never use Aspirin in children as it is associated with a dangerous condition called Reye's Syndrome
- Use only as directed by your physician
- You may give Motrin every 6 to 8 hours but not more than 4 times in a 24 hour period
- Keep all medication out of the reach of children
- Do not use in children under 6 months of age

Disclaimer: These guidelines are to help the caretaker with home treatment. If you are concerned about your child's health you should see your physician in person.