

UCLA OUTPATIENT REHABILITATION SERVICES	
<input type="checkbox"/> WESTWOOD 1000 Veteran Ave., A level Phone: (310) 794-1323 Fax: (310) 794-1457	<input type="checkbox"/> SANTA MONICA 1260 15 th St, Ste. 900 Phone: (310) 319-4646 Fax: (310) 319-2269
FOR APPTS, CALL: (310) 794-1323	
FAX: (310) 794-1457	

PATELLOFEMORAL INSTABILITY PHYSICAL THERAPY PRESCRIPTION

Diagnosis: (LEFT / RIGHT) Knee Patellofemoral Instability

Number of episodes: _____

Pathology: _____

Nonoperative: _____

Preoperative: _____

Progression of rehabilitation should be based upon the patient's ability to meet the goals of each phase. This rehabilitation protocol is designed for conservative management of a first-time acute patellar dislocation or for preoperative management of a patient planning to undergo surgical management in the near future.

PHASE I

Goals:

- ___ Decrease pain and swelling
- ___ Limit range of motion and weight-bearing to protect healing tissues
- ___ Return of muscle function
- ___ Avoid overaggressive therapy that may lead to patellofemoral pain syndrome

Suggested Intervention:

- ___ Bracing: set at 0° initially with ambulation, lateral buttress pad in place
- ___ Cryotherapy
- ___ Assess for McConnell taping
- ___ Instruction in partial weight-bearing activities with crutches
- ___ Electrical stimulation for activation of the VMO
- ___ Supine SLR with minimal to no pain
- ___ Ankle pumps for edema control
- ___ Isometric hamstrings

PHASE II

Criteria for advancement: No significant joint effusion, no quadriceps extension lag, minimal to no pain with ADLs

Goals:

- ___ Full ROM – pain free
- ___ Improve quadriceps strength
- ___ Low level functional activities
- ___ Initiate conditioning

Suggested Intervention:

- ___ Continue patellar bracing or taping
- ___ WBAT, d/c crutches when extension lag is no longer present
- ___ Continue electrical stimulation and modalities as needed
- ___ Continue supine SLR and add adduction/abduction SLRs
- ___ Toe raises with equal weightbearing
- ___ Closed kinetic chain exercises
- ___ Low level endurance and pool exercises

PHASE III

Criteria for advancement: Full active ROM, good to normal quadriceps strength, full weight-bearing with normal gait pattern

Goals:

- ___ Improve lower extremity/core function
- ___ Gradual return to high-level activities

Suggested Intervention:

- ___ Wean from bracing and taping as quadriceps function improves
- ___ Four-way hip exercises
- ___ Pool therapy – walking with progress to running
- ___ Sport and skill specific training
- ___ Proprioceptive training
- ___ Patient education

Criteria for Return to Full Activity: Equal ROM between lower extremities, no pain or edema, 85% strength compared with the uninjured limb, satisfactory 1-minute single leg hop test/two-legged hop test, patellar stability with clinical tests

Treatment: _____ times per week _____ Please institute home program for preoperative patients

Duration: _____ weeks

**Please send progress notes.

Physician's Signature: _____ Date: _____
Kristofer J. Jones, M.D., Attending Orthopaedic Surgeon

Patellofemoral Instability PAGE * MERGEFORMAT 2

Place label here

NAME OF PATIENT: _____

MRN: _____

Kristofer J. Jones, M.D.

Sports Medicine, Shoulder Surgery and Cartilage Restoration
UCLA Department of Orthopaedic Surgery
David Geffen School of Medicine at UCLA
10833 Le Conte Avenue, 76-143 CHS
Los Angeles, CA 90095-6902
Phone: (310) 825-6095
Fax: (310) 825-1311
CA License: A126262

Kristofer J. Jones, M.D.
UCLA Department of Orthopaedic Surgery