



## Colonoscopy Preparation using *Clenpiq* 2 Day PM/AM Split Dosing Regimen

The colon preparation is the most important step leading to the successful completion of a colonoscopy. We know it is inconvenient, but a clean colon will lead to an efficiently and safely performed colonoscopy that will lead to good gastrointestinal health. The cost of these FDA-approved preps range from \$30-\$100 depending on your insurance coverage, co-payment, and ability to use discount coupons. **Please pick up your prescription ahead of time or call your pharmacist to check your benefits and any cost savings card.** If still too expensive call our office. If you have chronic kidney disease, you will require a different preparation.

### **SEVEN DAYS BEFORE THE PROCEDURE**

#### **Medication restrictions – Please call the prescribing physician**

1. *Diabetics taking insulin*: will need dose adjustment the day before the exam. Please ask for advice from the physician that prescribes your insulin. For diabetics only taking diabetic pills simply stop the pills the day before the procedure (see below).
2. *Blood thinners (antithrombotic agents) including: aspirin, Coumadin (Warfarin), Plavix, Pradaxa, Xarelto, Eliquis, Brilinta and Effient*: These can increase the risk of bleeding during and after endoscopic procedures to a varying degree. Please ask the physician that prescribes the medication if, and when, you can stop these medications. **If** these medications are unable to be stopped you need to call our office.

#### **Items to Purchase**

1. Fill the prescription for the *Clenpiq*. (Pharmacy may have to order, can take up to 48 hours)
2. Fill the prescription for *Ondansetron* (anti-nausea medication).
3. Purchase two (2) Dulcolax laxative tablets (over the counter).
4. Any over-the-counter hemorrhoid ointment for anal discomfort (if needed).

#### **Diet restrictions**

No seeds, nuts, popcorn or peels from fruits and vegetables. Example: no peels from apples, potatoes, or cucumbers etc.

### **TWO DAYS BEFORE THE PROCEDURE**

1. Take two (2) Dulcolax tablets before going to bed.

### **ONE DAY BEFORE THE PROCEDURE**

#### **STARTING WHEN YOU WAKE UP, CLEAR LIQUID DIET ONLY FOR THE ENTIRE DAY.**

1. **You cannot have milk or orange juice for the entire day.**
2. Drink as much as you can to prevent dehydration. You may have bouillon, broth, black coffee (sugar and artificial sweeteners are okay, no creamer of any kind), tea (honey is okay), juice (apple, grape, cranberry), Gatorade, Powerade, lemonade, soda (regular or diet are okay), water, Popsicles and Jell-O (any flavor or color). Taking red Jell-O and red Gatorade may turn your bowel movements red and should be limited. If you have had dehydration before drink Pedialyte.
3. You may have clear candy (such as Lifesavers, Jolly Ranchers and gummy bears).
4. Do not take oral diabetic pills.

**TURN THE PAGE OVER FOR THE REST OF THE DIRECTIONS**

## **TIMING OF PREPARATION**

**Please follow these instructions and NOT the *CLENPIQ* box instructions**

### **EVENING BEFORE THE PROCEDURE**

This is a 2 step process divided into two sessions. You must drink the entire *CLENPIQ* to ensure your colon will be cleaned out. A poor prep can mean missed lesions (flat polyps) and cancellation and rescheduling of your procedure.

**Starting at 6:00pm (first dose, this CAN NOT be done earlier in the day.)**

1. Drink first bottle of CLENPIQ.
2. You will need to follow with at least 5 more 8 ounce glass of clear liquids, total of 40 ounces before bed.
3. **If you become nauseated, stop drinking the *CLENPIQ*. Take one of the *Ondansetron* tablets. Resume drinking the *CLENPIQ* after 30 minutes, or when the nausea passes. If you do not use the *Ondansetron* it can be kept in case you have nausea in the future.**
4. You can continue the clear liquid diet after you finish the 6:00pm dose of the prep.

### **THE MORNING OF THE PROCEDURE (second dose)**

**Beginning 4-6 hours prior to **scheduled procedure time**.** (Yes, this requires you to do early morning or in the middle of the night based on your procedure time.)

1. Drink second bottle of CLENPIQ.
2. You will need to follow with 4 more 8 ounce glass of clear liquids. You will need to have the additional 32 ounces of clear liquids drank at the latest 4 hours prior to your arrival time.
3. **If you become nauseated, stop drinking the *CLENPIQ*. Take one of the *Ondansetron* tablets. Resume drinking the *CLENPIQ* after 30 minutes, or when the nausea passes. If you do not use the *Ondansetron* it can be kept in case you have nausea in the future.**
4. **Take nothing by mouth, including water, after completing the last additional 24 ounces of clear liquids.**
5. You should take any necessary medications you normally use on the morning of the procedure with **sips** of water.
6. Arrive at your scheduled arrival time.
7. Bring a list of your medications, insurance cards and photo ID.
8. After the procedure you must have a responsible adult drive you home. It is best if your driver waits for you at the endoscopy center. **Taking a cab or Uber home is not an option.**
9. You may not drive or work until the next day.

### **IMPORTANT INFORMATION ABOUT SEDATION**

Sedatives used for endoscopy include Propofol. These medications provide amnesia and pain relief. You are breathing on your own.

### **QUESTIONS OR PROBLEMS:**

1. Daytime phone number is (314) 997-0554.
2. After hours number is (314) 388-6578.
3. Internet website address is [www.gidoctor.net](http://www.gidoctor.net)