

**PLEASE DISCONTINUE THE FOLLOWING MEDICATIONS AND/OR VITAMIN SUPPLEMENTS  
FOR THE STATED NUMBER OF DAYS PRIOR TO YOUR PAIN PROCEDURE**

Suspension Period (Days)	Generic Name	Brand Name(s)
14 Days (2 Weeks)	Ticlopidine	Ticlid
7 Days (1 Week)	Clopidogrel Bisulfate	Plavix
	Etanercept	Enbrel
	Mesalamine	Asacol, Rowasa
	Olsalazine Sodium	Dipentum
	Prasugrel	Effient
	Ticagrelor	Brillinta
5 Days	Abciximab	Reopro
	Aspirin (325 mg/day and above)	Bayer, Ecotrin, Durlaza
	Aspirin with Dipyridamole	Aggrenox
	Dabigatran	Pradaxa
	Warfarin	Coumadin
3 Days (72 Hours)	Apixaban	Eliquis
	Edoxaban	Savaysa
	Rivaroxaban	Xarelto
2 Days (48 Hours)	Cilostazol	Pletal
	Diclofenac	Arthrotec, Zipsor, Voltaren
	Fenoprofen	Fenortho, Nalfon
	Fondaparinux	Arixtra
	Etodolac	Lodine
	Ibuprofen	Advil, Motrin, Duexis
	Indomethacin	Indocin
	Ketorolac	Toradol, Acular
	Meloxicam	Mobic
	Nabumetone	Relafen
	Naproxen	Aleve, Naprosyn
1 Day (24 Hours)	Dalteparin	Fragmin
	Enoxaparin Sodium	Lovenox
	Eptifibatid	Integrilin
	Tirofiban	Aggrastat