



Surgical Consultants of Dallas, L.L.C

The name of your operation is robotic cholecystectomy with intra-operative cholangiogram which means that your gallbladder was removed and an x-ray was taken of your bile duct. Your surgery went well.

People live a normal life without a gallbladder. You do not have to change your diet or your lifestyle as a result of having it out. Some patients experience a slight increase in frequency of bowel movements for several weeks after this operation.

Follow-up Contacts

If you have questions about your condition, please call (972) 566-7860 to contact SURGICAL CONSULTANTS OF DALLAS. This number will be answered by our office during business hours and connect with an answering service at night and on weekends.

Specifically, call for any of the following:

- Your abdominal pain becomes much worse than when you left the hospital
- The area around your incisions becomes very red, increasingly tender, or begins to drain pus (Small amounts of blood-tinged fluid and mild redness are common and are no cause for concern.)
- Your temperature goes above 100.0° F (38° C)
- Persistent vomiting, diarrhea, or constipation develop
- You feel as if you are getting sick instead of well
- If you are considering contacting your other doctors or going to an emergency room for something related to the operation, please first call SURGICAL CONSULTANTS OF DALLAS (972) 566-7860. You will reach the office during business hours or be automatically forwarded to the on-call doctor if it is after hours.

Please phone (972) 566-7860 and make a follow-up appointment with Dr. Madni for 10-14 days following surgery. If no one answers, leave a message and your call will be returned within one business day.

Diet: You may eat a regular diet after surgery.

Pain Control and Other Medications:

When you leave the hospital, you will have a prescription for medicine to help with the pain from your incisions. Pain medications can cause constipation, so take a stool softener (Colace 200 mg twice daily) to avoid that while you are taking pain medications. You will also have a script for Zofran in case you experience nausea following surgery.



Surgical Consultants of Dallas, L.L.C

You may resume any medications you may have been taking for conditions unrelated to your gallbladder disease.

Bowel Movements:

Bowel movements may be irregular for several weeks, but they gradually return to normal. Pain medicine may make you constipated, and as the need for pain medicine decreases, so will the constipation. If you go for more than a couple of days without having a bowel movement, try eating prunes or taking a gentle laxative, such as milk of magnesia.

Wound Care:

Your incision is closed with absorbable sutures and surgical glue. You may take showers and allow soap and water to wash over incision. The glue will peel off in 10-14 days. Do not submerge your incision in water (no swimming or bathing) for one month following surgery.

Activity:

It is common in the first few days at home to feel a little more tired than expected. This comes from the increased activity involved in moving around home. It is not a problem, and is no reason to limit your activity. Just compensate by getting extra rest.

- Do not lift anything heavier than 15 pounds for two weeks following surgery.
- You may perform normal daily activities as you feel up to it. This includes walking up and down stairs, walking outside, traveling as a passenger in a car or a plane, etc.
- Do not drive while taking narcotic pain medications.