

# Kids & Teens Medical Group Vaccine Schedule 2021

<b>Birth</b>	
<b>HepB</b> <sup>1</sup>	

<b>Age 2 months</b>	
<b>DTaP</b> (Diphtheria, Tetanus, Pertussis)	
<b>Polio</b> (IPV)	
<b>HepB</b> <sup>2</sup> (age: 1-2 months)	1-2 months after birth dose
<b>Hib</b> (Hib meningitis)	
<b>PCV</b> (Pneumo)	
<b>RV</b> <sup>3</sup> (Rotavirus)	

<b>Age 4 months</b>	<b>Interval from previous dose</b>
<b>DTaP</b>	1-2 months
<b>Polio</b>	1-2 months
<b>HepB</b> <sup>2</sup>	1-2 months if 1st dose given at 2 months of age
<b>Hib</b>	1-2 months
<b>PCV</b>	1-2 months
<b>RV</b> <sup>3</sup>	4-10 weeks

<b>Age 6 months</b>	<b>Interval from previous dose</b>
<b>DTaP</b>	1-2 months
<b>Polio</b> (age: 6-18 months)	1-14 months
<b>HepB</b> <sup>2</sup> (age: 6-18 months)	2-12 months and at least 4 months after first dose
<b>Hib</b> <sup>4</sup>	1-2 months
<b>PCV</b>	1-2 months
<b>RV</b> <sup>3</sup>	4-10 weeks and if RV-5 (RotaTeq) used for doses 1 or 2

<b>Age 12 months</b>	<b>Interval from previous dose</b>
<b>HepA</b> (age: 12-23 months)	
<b>MMR</b> <sup>5,6</sup> (age: 12-15 months)	
<b>Varicella</b> <sup>6</sup> (age: 12-15 months)	
<b>Hib</b> (age: 12-15 months)	2-8 months
<b>PCV</b> <sup>7</sup> (age: 12-15 months)	6-8 weeks

<b>Age 15 months</b>	<b>Interval from previous dose</b>
<b>DTaP</b> <sup>8</sup>	6-12 months

<b>Age 18 months</b>	<b>Interval from previous dose</b>
<b>HepA</b>	6-18 months

<b>Age 4-6 years</b>	<b>Age 11-12 years</b>	<b>Age 16 years</b>
<b>DTaP</b>	<b>Tdap</b>	<b>Men ACWY (MCV4)</b>
<b>Polio (IPV)</b>	<b>HPV<sup>9</sup> (2 doses)</b>	<b>Men B<sup>10</sup></b>
<b>MMR<sup>5,6</sup></b>	<b>Men ACWY (MCV4)</b>	
<b>Varicella<sup>6</sup></b>		

**Every Fall: Flu Vaccine<sup>11</sup> Everyone 6 months and older**

**9 months - Check development and any missing vaccines**



(626) 298 7121 (ENGLISH)  
(626) 269 7744 (SPANISH)



CALL US  
(818) 361-5437