



Suburban Healthcare Associates

## Immunization, Lab and Developmental Screening Guidelines

We follow the Immunization Schedule as recommended by the American Academy of Pediatrics.

<b>NEWBORN</b>	<b>Hep B if not given in hospital</b>
<b>1 MONTH</b>	<b>Hep B #2 (must be 28 days apart)</b>
<b>2 MONTHS (post-natal depression screen)</b>	<b>DTaP, IPV, HIB(Pentacel), Prevnar, Rotateq</b>
<b>4 MONTHS</b>	<b>DTaP, IPV, HIB(Pentacel), Prevnar, Rotateq</b>
<b>6 MONTHS (post-natal depression screen)</b>	<b>DTaP, IPV, HIB(Pentacel), Prevnar, Rotateq</b>
<b>FLU</b>	<b>FLU</b>
<b>9 MONTHS (Chadis)</b>	<b>Hep B #3, Hgb, Lead (vaccine catch-up)</b>
<b>Ages &amp; Stages</b>	<b>Ages &amp; Stages</b>
<b>12 MONTHS (<u>must be 1 year old</u>)</b>	<b>Varivax, MMR, Prevnar, Hep A #1 (<u>must be 1 year old</u>)</b>
<b>15 MONTHS</b>	<b>DTaP, HIB, Hep A #1 (if not given at 12 mo)</b>
<b>18 MONTHS</b>	<b>Hep A #2 (15 mo vaccines if not given yet)</b>
<b>MCHAT &amp; Developmental</b>	<b>MCHAT &amp; Developmental</b>
<b>23 MONTHS</b>	<b>Hgb, Lead (15-18 mo vaccines if not given yet)</b>
<b>MCHAT</b>	<b>MCHAT</b>
<b>30 MONTHS</b>	<b>Vaccine catch-up, Ages and Stages</b>
<b>3 YEARS</b>	
<b>4 YEARS (<u>must be 4 years old</u>)</b>	<b>DTaP, IPV, MMR, Varicella (<u>must be 4 years old</u>)</b>
<b>5 YEARS</b>	<b>PPD &amp; 4 year vaccines if not given yet</b>
<b>6<sup>th</sup> GRADE</b>	<b>Tdap (booster every 10 years), Menactra (booster after age 16), HPV (series of 3)</b>

Please see next page for further information

**Yearly checkups are recommended by the American Academy of Pediatrics through age 21.**

**Flu shots are recommended for patients 6 months and older, generally given from August-June.**

**Fluoride treatments are recommended from age 9 months to 35 months up to 3 times per year.**

**If traveling, please call the office for recommended immunizations for the area you are traveling to.**