Alexander E. Weber, MD Sports Medicine Surgery Team Physician USC Athletics and LA Kings Appointments: 855-SC-SPORT Office: 818-658-5921 Cellular: 518-928-8389 Email: weberae@usc.edu Website: www.AlexWeberMD.com



ACI OF PATELLA / TROCHLEA WITHOUT AMZ REHABILITATION PROTOCOL

	WEIGHT BEARING	BRACE	ROM	EXERCISES
PHASE I 0-6 weeks	Full w/ brace	0-1 week: Locked in full extension at all times Off for CPM and exercise only 1-4 weeks:	0-6 weeks : Use CPM for 6 hours/day, beginning 0-30° for 0-2 weeks 2-4 weeks:	SLR, calf pumps, passive leg hangs to 45° at home
		Unlocked and worn daytime only Discontinue when	0-60° 4-6 weeks:	
		quads can control SLR w/o extension lag	0-90°	
PHASE II 6-8 weeks	Full	None	Full	Advance Phase I exercises
PHASE III 8- 12 weeks	Full	None	Full	Gait training, begin closed chain activities: wall sits, mini- squats, toe raises, stationary bike
				Begin unilateral stance activities, balance training
PHASE IV 12 weeks – 6 months	Full	None	Full	Advance Phase III exercises; maximize core/glutes, pelvic stability work, eccentric hamstrings
				May advance to elliptical, bike, pool as tolerated
PHASE V 6-	Full	None	Full	Advance functional activity
12 months				Return to sport-specific activity and impact when cleared by MD after 8 mos