



Botox®/Dysport®/Jeauveau®/Xeomin® Instructions

Pre-Treatment Instructions

- 7 DAYS BEFORE treatment (to prevent bruising): AVOID blood thinning over-the-counter medications such as Aspirin, Motrin, Ibuprofen, and Aleve. For mild pain or headaches Tylenol is preferred.
- Avoid herbal supplements, such as Garlic, Vitamin E, Ginkgo Biloba, St. John's Wort, and Omega-3 capsules.
- Do not drink alcoholic beverages 24 hours before (or after) your treatment to avoid extra bruising.
- Inform your provider if you have a history of Perioral Herpes to receive advice on antiviral therapy prior to treatment.
- Do not use Botulinum Toxins if you are pregnant or breastfeeding, are allergic to any of its ingredients, or suffer from any neurological disorders. Please inform your provider if you have any questions about this prior to the treatment.

Day of Treatment

- Arrive to the office with a "clean face". Please do not wear makeup. You may bring your own makeup to apply after your treatment.
- You will have photos taken prior to treatment. This is to evaluate and compare improvement of treatment at the time of your follow up appointment.
- You may experience small bumps on the injection sites for about 30 minutes post treatment
- You may experience a mild amount of tenderness or a stinging sensation following injection.
- Redness and swelling are normal. Some bruising may also be visible.
- You may experience some tenderness at the treatment site(s) that can last for a few hours or a few days. You may have bruises in the areas treated.

Immediately After Treatment

- Stay in a vertical position for 4 hours following injection. DO NOT "rest your head" or lie down; sit upright.
- You may apply an ice or cold gel pack to the area(s) treated (avoiding pressure) as this helps reduce swelling and the potential for bruising. You can apply the ice pack every hour for 20 minutes for 4 to 6 hours.
- Once you have adequately cooled/iced the area(s) as instructed and any pinpoint bleeding from the injection site(s) has subsided, you may begin wearing makeup.
- AVOID placing excessive pressure on the treated area(s) for the first few days; when cleansing your face or applying makeup, be very gentle.
- AVOID exercise or strenuous activities for the remainder of the treatment day; you may resume other normal activities/routines the day after treatment.
- You may take Acetaminophen/Tylenol if you experience any mild tenderness or discomfort.
- AVOID extended UV exposure until any redness/swelling has subsided. Be sure to apply an SPF 30 or higher sunscreen.
- If you have any questions or concerns, please contact the office.