



Liquid Nitrogen

Liquid nitrogen is applied to the top of the lesion, which may cause a stinging, burning pain that peaks about 2 mins after treatment. After the lesion and surrounding skin will become red and begin to swell. A blister may form within a few hours to days, and it can turn purple or black. This is not a cause of concern. The blister should flatten, and then the scab will slough off in 1-2 weeks.

Post care instructions: In most cases there is no care that is needed for these treated lesions besides washing daily with mild soap and water. Blisters provide a good natural bandage that protects the new skin growing underneath. If the roof of the blister opens or is removed, you should keep it bandaged with antibiotic ointment (Bacitracin or Polysporin) until it's healed. If the area is excessively uncomfortable, Tylenol will usually relieve it. The site should be protected from sun exposure while healing to prevent scar formation.



Liquid Nitrogen

Liquid nitrogen is applied to the top of the lesion, which may cause a stinging, burning pain that peaks about 2 mins after treatment. After the lesion and surrounding skin will become red and begin to swell. A blister may form within a few hours to days, and it can turn purple or black. This is not a cause of concern. The blister should flatten, and then the scab will slough off in 1-2 weeks.

Post care instructions: In most cases there is no care that is needed for these treated lesions besides washing daily with mild soap and water. Blisters provide a good natural bandage that protects the new skin growing underneath. If the roof of the blister opens or is removed, you should keep it bandaged with antibiotic ointment (Bacitracin or Polysporin) until it's healed. If the area is excessively uncomfortable, Tylenol will usually relieve it. The site should be protected from sun exposure while healing to prevent scar formation.



Liquid Nitrogen

Liquid nitrogen is applied to the top of the lesion, which may cause a stinging, burning pain that peaks about 2 mins after treatment. After the lesion and surrounding skin will become red and begin to swell. A blister may form within a few hours to days, and it can turn purple or black. This is not a cause of concern. The blister should flatten, and then the scab will slough off in 1-2 weeks.

Post care instructions: In most cases there is no care that is needed for these treated lesions besides washing daily with mild soap and water. Blisters provide a good natural bandage that protects the new skin growing underneath. If the roof of the blister opens or is removed, you should keep it bandaged with antibiotic ointment (Bacitracin or Polysporin) until it's healed. If the area is excessively uncomfortable, Tylenol will usually relieve it. The site should be protected from sun exposure while healing to prevent scar formation.



Liquid Nitrogen

Liquid nitrogen is applied to the top of the lesion, which may cause a stinging, burning pain that peaks about 2 mins after treatment. After the lesion and surrounding skin will become red and begin to swell. A blister may form within a few hours to days, and it can turn purple or black. This is not a cause of concern. The blister should flatten, and then the scab will slough off in 1-2 weeks.

Post care instructions: In most cases there is no care that is needed for these treated lesions besides washing daily with mild soap and water. Blisters provide a good natural bandage that protects the new skin growing underneath. If the roof of the blister opens or is removed, you should keep it bandaged with antibiotic ointment (Bacitracin or Polysporin) until it's healed. If the area is excessively uncomfortable, Tylenol will usually relieve it. The site should be protected from sun exposure while healing to prevent scar formation.