

CHAPTER-V. REFERENCES

Adams GA, Rose JK. Structural requirements of a membrane-spanning domain for protein anchoring and cell surface transport. *Cell* 1985; 41: 1007-1015.

Adams MD, Kerlavage AR, Fleischmair RD, Fuidner RA, Bult CJ, Lee NH, Kirkness EF, Weinstock KG, Gocayne JD, White O, Sutton G, Blake JA, Brandon RC, Chiu MW, Clayton RA, Cline RT, Cotton MD, Earle-Hughes J, Fine LD, FitzGerald LM, FitzHugh WM, Pritchman JL, Geoghagen NSM, Glodek A, Gnehm CL, Manna MC, Hedblom E, Hinkle PS, Kelley JM, Klimek ICM, Kelley JC, Liu LI, Marmaros SM, Merrick JM, Moreno-Palauques RF, McDonald LA, Nguyen DT, Pellegrino SM, Phillips CA, Ryder SE, Scott JL, Saudek DM, Shirley R, Small Ky, Spriggs TA, Utterback TR, Weidman JF, Li Y, Barthlow R, Bednarik DP, Cao L, Cepeda MA, Coleman TA, Collins EJ, Dimke D, Feng P, Ferrie A, Fischer C, Hastings GA, He WW, Hu iS, Huddleston KA, Greene JM, Gruber J, Hudson P, Kim A, Kozak DL, Kunsch C, Ji H, Li H, Meissner PS, Olsen H, Raymond L, Wei YF, Wing J, Xu C, Yu GL, Ruben SM, Dillon PJ, Fanrion MR, Rosen CA, Haseltine WA, Fields C, Fraser CM, Venter JC. Initial assessment of human gene diversity and expression patterns based upon 83 million nucleotides of cDNA sequence. *Nature* 1995; 377 (Suppl): 3-174 (see page-135).

Adams MD, Soares MB, Kerlavage AR, Fields C, Venter JC. Rapid cDNA sequencing (expressed sequence tags) from a directionally cloned human infant brain cDNA library. *Nature Genet* 1993; 4: 373-380.

Adams RD, Victor M. The neurology of aging. In: *Principles of Neurology*, 5th ed. McGraw-Hill, New York, 1993, pp 526-536.

Akhurst Ri, Lehnert SA, Gatherer D, Duffie E. The role of TGF β in mouse development. *Ann NY Acad Sci USA* 1990; 593: 259-271.

Akiyama S, Endo H, Inohara N, Ohta S, Kagawa Y. Gene structure and cell type-specific expression of the human ATP synthase a subunit. *Biochim Biophys Acta* 1994; 1219: 129-140.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol* 1990; 215: 403-410.

Alvarez IS, Schoenwolf GC. Expansion of surface epithelium provides the major intrinsic force for bending of the neural plate. *J Exp Zool* 1992; 261: 340-348.

Ang SL. The brain organization. *Nature* 1996; 380: 25-27.

Aota S, Ikemura T. Diversity in G + C content at the third position of codons in vertebrate genes and its cause. *Nuc Acids Res* 1986; 14: 6315-6355.

Asashima M, Nikano H, Shimada K, Kinoshita K, Ishii K, Shibai H, Veno N. Mesodermal induction in early amphibian embryos by activin A (erythroid differentiation factor). *Wilhelm Rota Arch Dev Biol* 1990; 198: 330-335.

- Ball MJ. Neuronal loss, neurofibrillary tangles and granulovacuolar degeneration in the hippocampus with aging and dementia. *Acta Neuropathol* 1977; 27: 111-118.
- Balvay L, Libri D, Fiszman MY. Pre-mRNA secondary structure and the regulation of splicing. *Bioessays* 1993; 15: 165-169.
- Basler K, Edlund T, Jessell TM, Yamada T. Control of cell pattern in the neural tube: Regulation of cell differentiation by dorsalin-1, a novel TGF β family member. *Cell* 1993; 73: 687-702.
- Benson D, Lipman DJ, Ostell J. GenBank. *Nuc Acids Res* 1993; 21: 2963-2965.
- Berget S. Exon recognition in vertebrate splicing. *J Biol Chem* 1995; 270: 2411-2414.
- Berry M, Rogers AW. The migration of neuroblasts in the developing cerebral cortex. *J Anat* 1965; 99: 691-709.
- Blasco MA, Funk W, Villeponteau B, Greider CW. Functional characterization and developmental regulation of mouse telomerase RNA. *Science* 1995; 269: 1267-1270.
- Bhat RV, Baraban JM, Johnson RC, Eipper BA, Mains RE. High levels of expression of the tumor suppressor gene APC during development of the rat central nervous system. *J Neurosci* 1994; 14: 3059-3071.
- Birgbauer E, Fraser SE. Violation of cell lineage restriction compartments in the chick hindbrain. *Development* 1994; 120: 1347-1356.
- Bopp D, Burn M, Baumgartner S, Frigerio G, Noll M. Conservation of a large protein domain in the segmentation gene *paired* and in functionally related genes in *Drosophila*. *Cell* 1986; 47: 1033-1040.
- Boyer PD. A perspective of the binding change mechanism for ATP synthesis. *FASEB J* 1989; 3: 2164-2178.
- Brandt CA, Kierkegaard O, Hindkjar J, Jensen PKA, Pedersen S. Ring chromosome 20 with loss of telomeric sequences detected by multicolor PRINS. *Clin Genet* 1993; 44: 26-31.
- Breathnach R, Chambon P. Organization and expression of eukaryotic split genes coding for proteins. *Annu Rev Biochem* 1981; 50: 349-383.
- Brown-Schimer S, Johnson KA, Lawrence JB, Johnson C, Bruskin A, Green NR, Hill DE. Molecular cloning and chromosome mapping of the human gene encoding protein phosphotyrosyl phosphatase IB. *Proc Natl Acad Sci USA* 1990; 87: 5148-5152.

Brunak S, Engelbrecht J, Knudsen S. Prediction of human mRNA donor and acceptor sites from the DNA sequence. *J Mol Biol* 1991; 220: 49-65.

Buell SJ, Coleman PD. Dendritic growth in the aged human brain and failure of growth in senile dementia. *Science* 1979; 206: 854-856.

Burnside B. Microtubules and microfilaments in newt neurulation. *Dev Biol* 1971; 26: 416-441.

Burnside B. Microtubules and microfilaments in amphibian neurulation. *Am Zool* 1973; 13: 989-1006.

Byrne E, Dennett X, Trounce I. Oxidative energy failure in post-mitotic cells: A major factor in senescence. *Revue Neurologique* 1991a; 147: 532-535.

Byrne E, Trounce I, Deimett X. Mitochondrial theory of senescence: Respiratory chain protein studies in human skeletal muscle. *Mech Ageing Dev* 1991b; 60: 295-302.

Campisi J. Replicative senescence: An old lives' tale? *Cell* 1996; 84: 497-500.

Carlin C, Phillips PD, Brooks-Frederich K, Knowles BB, Cristofalo VJ. Cleavage of epidermal growth factor receptor by a membrane-bound leupeptin-sensitive protease active in nonionic detergent lysates of senescent but not young human diploid fibroblasts. *J Cell Physiol* 1994; 160: 427-434.

Carpenter EM, Goddard JM, Chisaka O, Manley NR, Capecchi MR. Loss of Hox-A1 (Hox1.6) function results in the reorganization of the murine hindbrain. *Development* 1993; 118: 1063-1075.

Carroll SB. Homeotic genes and the evolution of arthropods and chordates. *Nature* 1995; 376: 479-485.

Catania J, Fairweather DS. DNA methylation and cellular aging. *Mutation Res* 1991; 256: 283-293.

Chang Z-F, Chen K-Y. Regulation of ornithine decarboxylase and other cell cycle-dependent genes during senescence of IMR-90 human diploid fibroblasts. *J Biol Chem* 1988; 263: 11431-11435.

Chapman T, Liddle LF, Kalb JM, Wolfner MF, Partridge L. Cost of mating in *Drosophila melanogaster* females is mediated by male accessory gland products. *Nature* 1995; 373: 241-244.

Chenn A, McConnell SK. Cleavage orientation and asymmetric inheritance of Notch immunoreactivity in mammalian neurogenesis. *Cell* 1995; 82: 631-641.

- Cho K W Y, Blumberg B, Steinbeisser II, De Robertis EM. Molecular nature of Spemann's organizer: The role of the *Xenopus* homeobox gene *gooseoid*. *Cell* 1991; 67: 1111-1120.
- Chomczynski P, Sacchi N. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 1987; 162: 156-159.
- Christian JL, Moon RT. Interactions between Xwnt-8 and Spemann organizer signalling pathways generate dorsoventral pattern in the embryonic mesoderm of *Xenopus*. *Genes Dev* 1993; 7: 13-28.
- Christy B, Nathans D. DNA binding site of the growth factor-inducible protein Zif268. *Proc Natl Acad Sci USA* 1989; 86: 8737-8741.
- Clark JM. Novel non-templated nucleotide addition reactions catalyzed by procaryotic and eucaryotic DNA polymerases. *Nuc Acids Res* 1988; 16: 9677-9686.
- Cohen BJ, Anver MR, Ringler DII, Adelman RC. Age-associated pathological changes in male rats. *Fed Proc* 1978; 37: 2848-2850.
- Coleman, GL, Barthold SW, Osbaldiston GW, Foster SJ, Jonas AM. Pathological changes during aging in barrier-reared Fischer 344 male rats. *J Gerontol* 1977; 32: 258-278.
- Collins K, Kobayashi R, Greider CW. Purification of *Tetrahymena* telomerase and cloning of genes encoding the two components of the enzyme. *Cell* 1995; 81: 677-686.
- Colman, PD, Kaplan BB, Osterburg HH, Finch CE. Brain poly(A)RNA during aging: Stability of yield and sequence complexity in two rat strains. *J Neurochem* 1980; 34: 335-345.
- Cooney CA. Are somatic cells inherently deficient in methylation metabolism? A proposed mechanism for DNA methylation loss, senescence and aging. *Growth Dev Aging* 1993; 57: 261-273.
- Costello B, Meymandi A, Freeman JA. Factors influencing gap-43 gene expression in PC12 pheochromocytoma cells. *J Neurosci* 1990; 10: 1398-1406.
- Cotter PD, Tumewu P, Browett PJ. Deletion of the long arm of chromosome 20 in a patient with small cell lymphocytic lymphoma. *Cancer Genet Cytogenet* 1993; 70: 142-143.
- Crook T, Bartus RT, Ferris SH. Age-associated memory impairment: Proposed diagnostic criteria and measures of clinical change - report of a National Institute of Mental Health Work Group. *Dev Neuropsychol* 1986; 2:261.
- Crossley PH, Martinez S, Martin GR. Midbrain development induced by FGF8 in the chick embryo. *Nature* 1996; 380: 66-68.

Crowley C, Curtis HJ. The development of somatic mutations in mice with age. *Proc Natl Acad Sci USA* 1963; 49: 626-628.

Cunliffe V, Smith JC. Ectopic mesoderm formation in *Xenopus* embryos caused by widespread expression of a brachyury homologue. *Nature* 1992; 358: 427-430.

Cunliffe V, Smith JC. Specification of mesodermal pattern in *Xenopus laevis* by interactions between Brachyury, noggin and Xwnt-8. *EMBO J* 1994; 13: 349-359.

Cutler RG. Transcription of unique and reiterated DNA sequences in mouse liver and brain tissues as a function of age. *Exp Gerontol* 1975; 10: 37-60.

D'Arcangelo G, Miao GG, Chen S-C, Soares HD, Morgan JI, Curran T. A protein related to extracellular matrix proteins deleted in the mouse mutant reeler. *Nature* 1995; 374: 719-723.

Davids M, Loppnow B, Tiedemann H. Neural differentiation of amphibian gastrula ectoderm exposed to phorbol ester. *Wilhelm Rota Arch Dev Biol* 1987; 196: 137-140.

Davis MP, Dewald GW, Pierre RV, Hoagland HC. Hematologic manifestations associated with deletions of the long arm of chromosome 20. *Cancer Genet Cytogenet* 1984; 12: 63-71.

Dell'Orco RT, Whittle WL. Nuclear matrix composition and in vitro cellular senescence. *Exp Gerontol* 1994; 29: 139-149.

De Robertis EM, Sasai Y. A common plan for dorsoventral patterning in Bilateria. *Nature* 1996; 380: 37-40.

Desmond ME, Field M. Evaluation of neural fold fusion and coincident initiation of spinal cord occlusion in chick embryo. *J Comp Neurol* 1992; 319: 246-260.

Desmond ME, Schoenwolf GC. Evaluation of the roles of intrinsic and extrinsic factors in occlusion of the spinal neurocoel during rapid brain enlargement in the chick embryo. *J Embryol Exp Morphol* 1986; 97: 25-46.

Dice JF. Altered degradation of proteins microinjected into senescent human fibroblasts. *J Biol Chem* 1982; 257: 14624-14627.

Dice JF. Cellular and molecular mechanisms of aging. *Physiol Rev* 1993; 73: 149-159.

Diez-Martin JL, Graham DL, Pettitt RM, Dewald GW. Chromosome studies in 104 patients with polycythemia vera. *Mayo Clin Proc* 1991; 66: 287-299.

DiPaolo BR, Pignolo RJ, Cristofalo VJ. Identification of proteins differentially expressed in quiescent and proliferatively senescent fibroblast cultures. *Exp Cell Res* 1995; 220: 178-185.

D'Mello NP, Jazwinski SM. Telomere length constancy during aging of *Saccharomyces cerevisiae*. *J Bacteriol* 1991; 173: 6709-6713.

Dolle P, Lufkin T, Krumlauf R, Mark M, Duboule D, Chambon P. Local alterations of Krox20 and Hox gene expression in the hindbrain suggest lack of rhombomeres 4 and 5 in homozygote null *Hoxa-1* (*Hox-1.6*) mutant embryos. *Proc Natl Acad Sci USA* 1993; 90: 7666-7670.

Dou D, Joseph R. Human Homolog of a novel mammalian brain developmental gene. Proceedings of the meeting *Human Genome 1994: Genes and Beyond*, Washington D.C. October 2-5, 1994 #24, p20 (abstract).

Dou D, Joseph R. Molecular cloning of human neuronatin. Proceedings of 47th meeting of the American Academy of Neurology, Seattle, WA. May 1995. *Neurology* 1995a; 45 (Suppl 4): A355 (abstract).

Dou D, Joseph R. Structure and expression of a novel human brain developmental gene. Proceedings of the 25th meeting of the Society for Neuroscience, San Diego, CA. November 1995. *Soc Neurosci Abs* 1995b; 21: 284 (abstract).

Dou D, Joseph R. Human *neuronatin* gene maps to chromosome 20q11.2-12. Proceedings of the 48th meeting of the American Academy of Neurology, San Francisco, CA. March 1996. *Neurology* 1996a; 46: A148 (abstract).

Dou D, Joseph R. The structure and organization of the human *neuronatin* gene. *Genomics* 1996b; 33: 292-297.

Dou D, Joseph R. Cloning of human *neuronatin* gene and its localization to chromosome-20q11.2-12: The deduced protein is a novel "proteolipid". *Brain Res* 1996c; 723: 8-22.

Dykhuisen D. The evolution of cell senescence, atherosclerosis, and benign tumours. *Nature* 1974; 251: 616-618.

Echelard Y, Epstein DJ, St-Jacques B, Shen L, Mohler J, McMahon JA, McMahon AP. Sonic hedgehog, a member of a family of putative signaling molecules, is implicated in the regulation of CNS polarity. *Cell* 1993; 75: 1417-1430.

Ekker SC, Jackson DG, von Kessler DP, Sun BI, Young KE, Beachy PA. The degree of variation in DNA sequence recognition among four *EMBO J* 1994; 13: 3551-3560.

Eldred GE, Lasky MR. Retinal age pigments generated by self-assembling lysosomotropic detergents. *Nature* 1993; 361: 724-726.

Epstein DJ, Vekemans M, Gros P. *splotch* (SP^{2H}), a mutation affecting development of the

mouse neural tube, shows a deletion within the paired homeodomain of *Pax-3*. *Cell* 1991; 67: 767-774.

Erickson CA, Tosney KW, Weston JA. Analysis of migrating behaviour of neural crest and fibroblastic cells in embryonic tissues. *Dev Biol* 1980; 77: 142-156.

Erickson CA, Weston JA. An SEM analysis of neural crest migration in the mouse. *J Embryol Exp Morphol* 1983; 74: 97-118.

Erickson-Viitanen S, Ruggieri S, Natalini P, Horecker BL. Thymosin β_{10} , a new analog of thymosin β_4 in mammalian tissues. *Arch Biochem Biophys* 1983; 225: 407-413.

Ericson J, Muhr J, Placzek M, Lints T, Jessell TM, Edlund T. Sonic hedgehog induces the differentiation of ventral forebrain neurons: A common signal for ventral patterning within the neural tube. *Cell* 1995; 81: 747-756.

Ericson J, Thor S, Edlund T, Jessell TM, Yamada T. Early stages of motor neuron differentiation revealed by expression of homeobox gene *Islet-1*. *Science* 1992; 256: 1555-1560.

Faust JR, Rodman JS, Daniel PF, Dice JF, Bronson RT. Two proteolipids and dolichol-linked oligosaccharides accumulate in motor neuron degeneration mice (*mnd/mnd*), a model for neuronal ceroid lipofuscinosis. *J Biol Chem* 1994; 269: 10150-10155.

Fearnley IM, Walker JE, Martinus RD, Jolly RD, Kirkland KB, Shaw GJ, Palmer DN. The sequence of the major protein stored in ovine ceroid lipofuscinosis is identical with that of the dicyclohexylcarbodiimide-reactive proteolipid of mitochondrial ATP synthase. *Biochem J* 1990; 268: 751-758.

Feng J, Funk WD, Wang S-S, Weinrich SL, Avilion AA, Chiu C-P, Adams RR, Chang E, Allsopp RC, Yu J, Le S, West MD, Harley CB, Andrews WH, Greider CW, Villeponteau B. The RNA component of human telomerase. *Science* 1995; 269: 1236-1241.

Finch CE. In: *Longevity, Senescence and the Genome*. University of Chicago Press, 1990.

Finch CE. Neuron atrophy during aging: programmed or sporadic. *Trends Neurosci* 1993a; 16: 104-110.

Finch CE. Biochemistry of aging in the brain. In: *Basic Neurochemistry*, 5th ed, Siegel GJ, Agranoff BW, Albers RW, Molinoff PB, eds. Raven press, New York, 1993b.

Fishell G. Striatal precursors adopt cortical identities in response to local cues. *Development* 1995; 121: 803-812.

Fishell G, Mason CA, Hatten ME. Dispersion of neural progenitors within the germinal zones

of the forebrain. *Nature* 1993; 362: 636-638.

Fornace AJ, Alamo I, Hollander MC. DNA damage-inducible transcripts in mammalian cells. *Proc Natl Acad Sci USA* 1988; 85: 8800-8804.

Fornace AJ, Mitchell JB. Induction of B2 RNA polymerase III transcription by heat shock: Enrichment for heat shock induced sequences in rodent cells by hybridization subtraction. *Nuc Acids Res* 1986; 14: 5793-5811.

Frantz GD, Bohner AP, Akers RM, McConnell SK. Regulation of POU domain gene SCIP during cerebral cortical development. *J Neurosci* 1994a; 14: 472-485.

Frantz GD, Weimann JM, Levin ME, McConnell SK. Otx1 and Otx2 define layers and regions in developing cerebral cortex and cerebellum. *J Neurosci* 1994b; 14: 5725-5740.

Fraser S, Keynes R, Lumsden A. Segmentation in the chick embryo hindbrain is defined by cell lineage restrictions. *Nature* 1990; 344: 431-435.

Frederich KB, Phillips PD, Cristofalo VJ. Stimulation of DNA synthesis in senescent human cells following incubation with plasma membranes. *Exp Cell Res* 1992; 202: 386-390.

Fujita, S. Kinetics of cellular proliferation. *Exp Cell Res* 1962; 28: 52-60.

Fujita S. Analysis of neuron differentiation in the central nervous system by tritiated thymidine autoradiography. *J Comp Neurol* 1964; 122: 311-328.

Fujita S. Application of light and electron microscopy to the study of the cytogenesis of the forebrain. In: *Evolution of the Forebrain*, Hassler R, Stephen H, eds. Plenum, New York, 1966, pp 180-196.

Fujita T, Mandel JL, Shirasawa T, Hino O, Shirai T, Maruyama N. Isolation of cDNA clone encoding human homologue of senescence marker protein-30 (SMP30) and its location on the X chromosome. *Biochem Biophys Acta* 1995; 1263: 249-252.

Galileo DS, Gray GE, Owens GC, Majors J, Sanes JR. Neurons and glia arise from a common progenitor in chick optic tectum: Demonstration with two retroviruses and cell type-specific antibodies. *Proc Natl Acad Sci USA* 1990; 87: 458-462.

Gehring WJ, Qian YQ, Billeter M, Furukubo-Tokunaga K, Schier AF, Resendez-Perez D, Affolter M, Otting G, Wuthrich K. Homeodomain-DNA recognition. *Cell* 1994; 78: 211-223.

Gems D, Riddle DL. Longevity in *Caenorhabditis elegans* reduced by mating but not gamete production. *Nature* 1996; 379: 723-725.

- Gilbert SF. In: *Developmental Biology*, 4th ed. Sinauer Associates Inc., Sunderland, MA, 1994.
- Gilley D, Balckburn EH. Lack of telomere shortening during senescence in *Paramecium*. *Proc Natl Acad Sci USA* 1994; 91: 1955-1958.
- Goffinet AM. A real gene for reeler. *Nature* 1995; 374: 675-676.
- Golden JA, Chernoff GF. Intermittent pattern of neural tube closure in two strains of mice. *Teratology* 1993; 47: 73-80.
- Goldman J, Cote L. In: *Principles of Neural Science*, 3rd ed, Kandel ER, Schwartz JH, Jessell TM, eds. Appleton and Lange, Norwalk, 1991.
- Goldspink DF, Kelly FJ. Protein turnover and growth in the whole body, liver, and kidney of the rat from foetus to senility. *Biochem J* 1984; 217: 507-516.
- Goldstein S, Moerman EJ, Baxter RC. Accumulation of insulin-like growth factor binding protein-3 in conditioned medium of human fibroblasts increases with chronologic age of donor and senescence in vitro. *J Cell Physiol* 1993; 156: 294-302.
- Gompertz B. On the nature of the function expressive of the law of human mortality and a new mode of determining the value of life contingencies. *Phil Trans Roy Soc Lond* 1825; 115: 513-585.
- Goodman L, Stein GH. Basal and induced amounts of interleukin-6 mRNA decline progressively with age in human fibroblasts. *J Biol Chem* 1994; 269: 19250-19255.
- Gottlieb S, Ruvkun G. *daf-2*, *daf-16* and *daf-23*: Genetically interacting genes controlling dauer formation in *Caenorhabditis elegans*. *Genetics* 1994; 137: 107-120.
- Goulding MD, Chalepakis G, Deutsch U, Erselius J, Gruss P. Pax-3, a novel murine DNA binding protein expressed during early embryogenesis. *EMBO J* 1991; 10: 1135-1147.
- Green JBA. Roads to neuralness: Embryonic neural induction as derepression of a default state. *Cell* 1994; 77: 317-320.
- Green JBA, New HV, Smith JC. Responses of embryonic *Xenopus* cells to activin and FGF are separated by multiple dose thresholds and correspond to distinct axes in the mesoderm. *Cell* 1992; 71: 731-739.
- Green JBA, Smith JC. Graded changes in dose of a *Xenopus* activin A homologue elicit stepwise transitions in embryonic cell fate. *Nature* 1990; 347: 391-394.

Green JBA, Smith JC, Gerhart JC. Slow emergence of a multithreshold response to activin requires cell-contact dependent sharpening but not prepattern. *Development* 1994; 120: 2271-2278.

Green MR. Biochemical mechanisms of constitutive and regulated pre-mRNA splicing. *Annu Rev Cell Biol* 1991; 7: 559-599.

Greene LA, Tischler AS. Establishment of a noradrenergic clonal line of rat adrenal pheochromocytoma cells which respond to nerve growth factor. *Proc Natl Acad Sci USA* 1976; 73: 2424-2428.

Grigoriev VG, Moerman EJ, Goldstein S. Senescence and cell density of human diploid fibroblasts influence metabolism of insulin-like growth factor binding proteins. *J Cell Physiol* 1994; 160: 203-211.

Grove EA, Williams BP, Li DQ, Hajihosseini M, Friedrich A, Price J. Multiple restricted lineages in the embryonic rat cerebral cortex. *Development* 1993; 117: 553-561.

Gruss P, Walther C. Pax in development. *Cell* 1992; 69: 719-722.

Gualandi F, Morelli C, Pavan JV, Rimessi P, Sensi A, Bonfatti A, Gruppioni R, Possati L, Stanbridge EJ, Barbanti-Brodano G. Induction of senescence and control of tumorigenicity in BK virus transformed mouse cells by human chromosome 6. *Genes Chromosomes Cancer* 1994; 10: 77-84.

Guerrieri F, Capozza G, Kalous M, Papa S. Age-related changes of mitochondrial F₀F₁ ATP synthase. *Ann NY Acad Sci* 1992; 671: 395-402.

Gurdon JB, Mitchell A, Mahony D. Direct and continuous assessment by cells of their position in a morphogen gradient. *Nature* 1995; 376: 520-521.

Guthrie S, Lumsden A. Formation and regeneration of rhombomere boundaries in the developing chick hindbrain. *Development* 1991; 112: 221-229.

Guthrie S, Price V, Lumsden A. Selective dispersal of avian rhombomere cells in orthotopic and heterotopic grafts. *Development* 1993; 118: 527-538.

Hall AK, Chen S-C, Hempstead JL, Morgan JI. Retinoic acid regulates thymosin β 10 levels in rat neuroblastoma cells. *J Neurochem* 1991; 56: 462-468.

Hamburger V. Historical landmarks in neurogenesis. *Trends Neurosci* 1981; 4: 151-155.

Harley CB. Telomere loss: Mitotic clock or genetic time bomb? *Mutation Res* 1991; 256: 271-282.

Harley CB, Futcher AB, Greider CW. Telomeres shorten during ageing of human fibroblasts. *Nature* 1990; 345: 458-460.

Hawkins JD. A survey on intron and exon lengths. *Nuc Acids Res* 1988; 16: 9893-9908.

Hayflick L. The cell biology of human aging. *N Engl J Med* 1976; 295: 1302-1308.

Hayflick L, Moorhead PS. The serial cultivation of human diploid cell strains. *Exp Cell Res* 1961; 25: 585-621.

He X, Treacy MN, Simmons DM, Ingraham HA, Swanson LW, Rosenfeld MG. Expression of large family of POU-domain regulatory genes in mammalian brain development. *Nature* 1989; 340: 35-42.

Hemmati-Brivanlou A, Kelly OG, Melton DA. Follistatin, an antagonist of activin, is expressed in the Spemann organizer and displays direct neuralising activity. *Cell* 1994; 77: 283-295.

Hemmati-Brivanlou A, Melton DA. A truncated activin receptor inhibits mesoderm induction and formation of axial structures in *Xenopus* embryos. *Nature* 1992; 359: 609-614.

Hendricks-Taylor LR, Bachinski LL, Siciliano MJ, Fertitta A, Trask B, deJong PJ, Ledbetter DH, Darlington GJ. The CCAAT enhancer binding protein (c/EBP α) gene (CEBPA2) maps to human chromosome 19q19.1 and the related nuclear factor NF-IL6 (c/EBP β gene C/EBPB) maps to human chromosome 20q13.1. *Genomics* 1992; 14: 12-17.

Hensler PJ, Annab LA, Barrett JC, Pereira-Smith OM. A gene involved in control of human cellular senescence on human chromosome 1q. *Mol Cell Biol* 1994; 14: 2291-2297.

Herr W, Sturm RA, Clerc RG, Corcoran LM, Baltimore D, Sharp PA, Ingraham HA, Rosenfeld MG, Finney M, Ruvkun G, Horvitz HR. The POU domain: A large conserved region in mammalian *pit-1*, *oct-1*, *oct-2* and *Caenorhabditis elegans unc-86* gene products. *Genes Dev* 1988; 2: 1513-1516.

Hirsch M-R, Gaugler L, Deagostini-Bazin H, Bally-Cuif L, Goridis C. Identification of positive and negative regulatory elements governing cell-type-specific expression of the neural cell adhesion molecule gene. *Mol Cell Biol* 1990; 10: 1959-1968.

His W. Zur Geschichte des menschlichen Rückenmarks und der Nervenwurzeln. *Ges Wissensch BD* 1886; 13(5): 477.

His, W. Die Neuroblasten und deren entstehung im embryonalen Mark. *Arch Anat Entw Jg* 1889; 249-300.

Holley SA, Jackson PD, Sasai Y, Lu B, De Robertis EM, Hoffmann FM, Ferguson EL. A

- conserved system for dorsal-ventral patterning in insects and vertebrates involving sog and chordin. *Nature* 1995; 376: 249-253.
- Holt CE, Bertsch TW, Ellis HM, Harris WA. Cellular determination in the *Xenopus* retina is independent of lineage and birthdate. *Neuron* 1988; 1: 15-26.
- Hubbard K, Dhanaraj SN, Sethi KA, Rhodes J, Wilusz J, Small MB, Ozer HL. Alteration of DNA and RNA binding activity of human telomere binding proteins occurs during cellular senescence. *Exp Cell Res* 1995; 218: 241-247.
- Huff K, End D, Guroff G. Nerve growth factor-induced alteration in the response of PC12 pheochromocytoma cells to epidermal growth factor. *J Cell Biol* 1981; 88: 189-198.
- Hughes KA, Charlesworth B. A genetic analysis of senescence in *Drosophila*. *Nature* 1994; 367: 64-66.
- Hunziker W. The 28-kDa vitamin D-dependent calcium-binding protein has a six domain structure. *Proc Natl Acad Sci USA* 1986; 83: 7578-7582.
- Hunziker W, Schrickel S. Rat brain calbindin D28: Six domain structure and extensive amino acid homology with chicken calbindin D28k. *Mol Endocrinol* 1988; 2: 465-473.
- Ekemura T. Codon usage and tRNA content in unicellular and multicellular organisms. *Mol Biol Evol* 1985; 2: 13-34.
- Izquierdo JM, Luis AM, Cuezva JM. Postnatal mitochondrial differentiation in rat liver. *J Biol Chem* 1990; 265: 9090-9097.
- Jacobs SM, Gorse KM, Westin EH. Identification of a second promoter in the human *c-myc* proto-oncogene. *Oncogene* 1994; 9: 227-235.
- Jacobson AG. Morphogenesis of the neural plate. In: *Morphogenesis and Pattern Formation*, Connely IG, et al. Raven Press, New York, 1981, pp 233-263.
- Jazwinski SM. Longevity, genes, and aging. *Science* 1996; 273: 54-59.
- Jazwinski SM, Chen JB, Sun J. A single gene change can extend yeast life span: The role of ras in cellular senescence. *Adv Exp Med Biol* 1993; 330: 45-53.
- Jendrisak J, Young RA, Engel JD. Cloning cDNA into λ gt10 and λ gt11. *Meth Enzymol* 1987; 152: 359-371.
- Joseph JA. The putative role of free radicals in the loss of neuronal functioning in senescence. *Integrative Physiological Behavioral Sci* 1992; 27: 216-227.

Joseph R, Dou D, Tsang W. Molecular cloning of a novel mRNA (Neuronatin) that is highly expressed in neonatal mammalian brain. *Biochem Biophys Res Commun* 1994; 201: 1227-1234.

Joseph R, Dou D, Tsang W. Neuronatin mRNA: Alternatively spliced forms of a novel brain-specific mammalian developmental gene. *Brain Res* 1995a; 690: 92-98.

Joseph R, Han E. Amyloid β -protein fragment 25-35 causes activation of cytoplasmic calcium in neurons. *Biochem Biophys Res Commun* 1992; 184: 1441-1447.

Joseph R, Tsang W. Molecular cloning of differentially expressed genes in brain aging. Proceedings of the 119th meeting of the American Neurological Association, San Francisco, CA. October 1994. *Ann Neurol* 1994; 36: 311 (abstract).

Joseph R, Tsang W, Dou D. The p23 transplantation antigen mRNA is differentially expressed in human fetal brain. *J Neural Sci* 1995b; 130: 224-227.

Joseph R, Tsang W, Han E, Saed GM. Neuronal β -amyloid precursor protein gene expression: Regulation by aurintricarboxylic acid. *Brain Res* 1993; 625: 244-255.

Kaplan RM. Imagine no coronary artery disease. *Circulation* 1991; 83: 1452-1455.

Karfunkel P. The activity of microtubules and microfilaments in neurulation in the chick. *J Exp Zool* 1972; 181: 289-302.

Kawasaki ES. Amplification of RNA. In: *PCR Protocols: A Guide of Methods and Application*, Innis MA, Gelfand DH, Sninsky JJ, White TJ, eds. Academic Press, Inc., San Diego, CA. 1990 pp 21-27.

Kennedy BK, Austriaco NR Jr, Zhang J, Guarente L. Mutation in the silencing gene SIR4 can delay aging in *S. cerevisiae*. *Cell* 1995; 80: 485-496.

Kenyon C. Ponce d'elegans: Genetic quest for the fountain of youth. *Cell* 1996; 84: 501-504.

Kenyon C, Chang J, Gensch E, Rudner A, Tabtiang R. A *C. elegans* mutant that lives twice as long as wild type. *Nature* 1993; 366: 461-464.

Kessel M, Gruss P. Murine developmental control genes. *Science* 1990; 249: 374-379.

Kessler DS, Melton DA. Vertebrate embryonic induction: Mesodermal and neural patterning. *Science* 1994; 266: 596-604.

Khan AS, Wilcox AS, Hopkins JA, Sikela JM. Efficient double stranded sequencing of cDNA clones containing long poly(A) tails using anchored poly(dT) primers. *Nuc Acids Res* 1991; 19: 1715.

- Kim JH, Alfieri AA, Rosenblum M, Bravo S, Kim SH. Low dose rate radiotherapy for transplantable gliosarcoma in the rat brain. *J Neuro Oncol* 1990; 9: 9-15.
- Klass MR. A method for the isolation of longevity mutants in the nematode *Caenorhabditis elegans* and initial results. *Mech Ageing Dev* 1983; 22: 279-286.
- Klee WA, Nirenberg M. A neuroblastoma x glioma hybrid cell line with morphine receptors. *Proc Natl Acad Sci USA* 1974; 71: 3474-3477.
- Komuro H, Rakic P. Selective role of N-type calcium channels in neuronal migration. *Science* 1992; 257: 806-809.
- Komuro H, Rakic P. Modulation of neuronal migration by NMDA receptors. *Science* 1993; 260: 95-97.
- Kordon EC, McKnight RA, Jhappan C, Hennighausen L, Merlino G, Smith GH. Ectopic TGF β 1 expression in the secretory mammary epithelium induces early senescence of the epithelial stem cell population. *Dev Biol* 1995; 168: 47-61.
- Kozak M. Structural features in eukaryotic mRNAs that modulate the initiation of translation. *J Biol Chem* 1991; 266: 19867-19870.
- Kraner SD, Chong JA, Tsay H-J, Mandel G. Silencing the type II sodium channel gene: A model for neural-specific gene regulation. *Neuron* 1992; 9: 37-44.
- Krueger BK, Burne JF, Raff MC. Evidence for large-scale astrocyte death in the developing cerebellum. *J Neurosci* 1995; 15: 3366-3374.
- Krumlauf R. Hox genes in vertebrate development. *Cell* 1994; 78: 191-201.
- Kumar S, Matsuzaki T, Yoshida Y, Noda M. Molecular cloning and biological activity of a novel developmentally regulated gene encoding a protein with β -transducin-like structure. *J Biol Chem* 1994; 269: 11318-11326.
- Kumar S, Tomooka Y, Noda M. Identification of a set of genes with developmentally down-regulated expression in the mouse brain. *Biochem Biophys Res Commun* 1992; 185: 1155-1161.
- Kunze N, Yang GC, Jiang ZY, Hameister H, Adolph S, Wiedorn KH, Richter A, Knippers R. Localization of the active type I DNA topoisomerase gene in human chromosome 20q11.2-13.1 and two pseudogenes on chromosomes 1q23-23 and 22q11.2-13.1. *Hum Genet* 1989; 84: 6-10.
- Kyte J, Doolittle RF. A simple method for displaying the hydropathic character of protein. *J Mol Biol* 1982; 157: 105-132.

Lahiri DK, Robakis NK. The promoter activity of the gene encoding Alzheimer β -amyloid precursor protein (APP) is regulated by blocks of upstream sequences. *Mol Brain Res* 1991; 9: 253-257.

Lamb TM, Knecht AK, Smith WC, Stachel SE, Ecomides AN, Stahl N, Yancopoulos GD, Harland RM. Neural induction by the secreted polypeptide noggin. *Science* 1993; 262: 713-718.

Langman J. Histogenesis of the central nervous system. In: *The Structure and Function of the Nervous System*, Bourne GH, ed. Academic Press, New York, 1968, pp 33-65.

Langman J, Guerrant RL, Freeman BG. Behavior of neuro-epithelial cells during closure of the neural tube. *J Comp Neurol* 1966; 127: 399-411.

Larsen PL, Albert PS, Riddle DL. Genes that regulate both development and longevity in *Caenorhabditis elegans*. *Genetics* 1995; 139: 1567-1583.

Leber SM, Breedlove SM, Sanes JR. Lineage, arrangement and death of clonally related motoneurons in chick spinal cord. *J Neurosci* 1990; 10: 2451-2462.

Lee JE, Hollenberg SM, Snider L, Turner DL, Lipnick N, Weintraub H. Conversion of *Xenopus* ectoderm into neurons by NeuroD, a basic helix-loop-helix protein. *Science* 1995; 268: 836-844.

Levitt P, Cooper ML, Rakic P. Coexistence of neuronal and glial precursor cells in the cerebral ventricular zone of the fetal monkey: An ultrastructural immunoperoxidase analysis. *J Neurosci* 1981; 1: 27-39.

Lewin B. The apparatus for nuclear splicing. In: *Genes V*, Lewin B, ed. Oxford University Press, Oxford, 1994, pp 911-940.

Li L, Suzuki T, Mori N, Greengard P. Identification of a functional silencer element involved in neuron-specific expression of the synapsin-I gene. *Proc Natl Acad Sci USA* 1993; 90: 1460-1464.

Liang P, Averboukh L, Keyomarsi K, Sager R, Pardee AB. Differential display and cloning of messenger RNAs from human breast cancer versus mammary epithelial cells. *Cancer Res* 1992; 52: 6966-6968.

Liang P, Averboukh L, Pardee AB. Distribution and cloning of eukaryotic mRNAs by means of differential display: Refinements and optimization. *Nuc Acids Res* 1993; 21: 3269-3275.

Liang P, Pardee AB. Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction. *Science* 1992; 257: 967-971.

- Lin S-C, Morrison-Bogorad M. Cloning and characterization of a testis-specific thymosin β 10 cDNA. *J Biol Chem* 1991; 266: 23347-23353.
- Liu Z, Lee A, Gilbert W. Gene disruption of a G4-DNA-dependent nuclease in yeast leads to cellular senescence and telomere shortening. *Proc Natl Acad Sci USA* 1995; 92: 6002-6006.
- Lois C, Garcia-Verdugo J-M, Alvarez-Buylla A. Chain migration of neuronal precursors. *Science* 1996; 271: 978-981.
- Lomri N, Perret C, Gouhier N, Thomasset M. Cloning and analysis of calbindin-d28k cDNA and its expression in the central nervous system. *Gene* 1989; 80: 87-98.
- London C, Akers R, Phillips C. Expression of Epi-1, an epidermis-specific marker in *Xenopus laevis* embryos, is specified prior to gastrulation. *Dev Biol* 1988; 129: 380-389.
- Lo Turco JJ, Kriegstein AR. Clusters of coupled neuroblasts in embryonic neocortex. *Science* 1991; 252: 563-566.
- Lugo DI, Chen S-C, Hall AK, Ziai R, Hempstead JL, Morgan JI. Developmental regulation of β -thymosins in the rat central nervous system. *J Neurochem* 1991; 56: 457-461.
- Lumpkin Jr CK, McClung K, Pereira-Smith OM, Smith JR. Existence of high abundance antiproliferative mRNA's in senescent human diploid fibroblasts. *Science* 1986; 232: 393-395.
- Lumsden A, Clarke JD, Keynes R, Fraser S. Early phenotypic choices by neuronal precursors, revealed by clonal analysis of the chick embryo hindbrain. *Development* 1994; 120: 1581-1589.
- Lumsden A, Keynes R. Segmental patterns of neuronal development in the chick hindbrain. *Nature* 1989; 337: 424-428.
- Lundblad V, Blackburn EH. An alternative pathway for yeast telomere maintenance rescues est1-senescence. *Cell* 1993; 73: 347-360.
- Luskin MB, Pearlman AL, Sanes JR. Cell lineage in the cerebral cortex of the mouse studied *in vivo* and *in vitro* with a recombinant retrovirus. *Neuron* 1988; 1: 635-647.
- Macieira-Coelho A. Chromatin reorganization during senescence of proliferating cells. *Mutation Res* 1991; 256: 81-104.
- Makino R, Akiyama K, Yasuda J, Mashiyama S, Honda S, Sekiya T, Hyashi K. Cloning and characterization of a c-myc intron binding protein (MIBP1). *Nuc Acids Res* 1994; 22: 5679-5685.
- Mandel M, Moriyama Y, Hulmes JD, Pan Y-CE, Nelson H, Nelson N. cDNA sequence

encoding the 16-kDa proteolipid of chromaffin granules implies gene duplication in the evolution of H⁺-ATPases. *Proc Natl Acad Sci USA* 1988; 85: 5521-5524.

Maniatis T, Goodbourn S, Fischer JA. Regulation of inducible and tissue-specific gene expression. *Science* 1987; 236: 1237-1245.

Mantell LL, Greider CW. Telomerase activity in germline and embryonic cells of *Xenopus*. *EMBO J* 1994; 13: 3211-3217.

Mark M, Lufkin T, Vonesch JL, Ruberte E, Olivo JC, Dolle P, Gorry P, Lumsden A, Chambon P. Two rhombomeres are altered in *Hoxa-1* mutant mice. *Development* 1993; 119: 319-338.

Marshall H, Nonchev S, Sham MH, Muchamore I, Lumsden A, Krumlauf R. Retinoic acid alters hindbrain Hox code and induces transformation of rhombomeres 2/3 into a 4/5 identity. *Nature* 1992; 360: 737-741.

Martin M, Nabout RE, Lafuma C, Crechet F, Remy J. Fibronectin and collagen gene expression during in vitro ageing of pig skin fibroblasts. *Exp Cell Res* 1990; 191: 8-13.

McConnell SK. Development and decision-making in the mammalian cerebral cortex. *Brain Res Bull* 1988; 13: 1-23.

McConnell SK. Strategies for the generation of neuronal diversity in the developing central nervous system. *J Neurosci* 1995; 15: 6987-6998.

McCormick A, Campisi J. Cellular aging and senescence. *Curr Opin Cell Biol* 1991; 3: 230-234.

McMahon AP, Bradley A. The *Wnt-1 (int-1)* proto-oncogene is required for development of a large region of the mouse brain. *Cell* 1990; 62: 1073-1085.

McMahon AP, Joyner AL, Bradley A, McMahon JA. The midbrain-hindbrain phenotype of *wnt-1/wnt-1* mice results from stepwise deletion of *engrailed-expressing* cells by 9.5 days postcoitum. *Cell* 1992; 69: 581-595.

McMahon AP, Moon RT. Ectopic expression of the proto-oncogene *int-1* in *Xenopus* embryos leads to duplication of the embryonic axis. *Cell* 1989; 58: 1075-1084.

Medrano EE, Yang F, Boissy R, Farooqui J, Shah V, Matsumoto K, Nordlund JJ, Park HY. Terminal differentiation and senescence in the human melanocyte: Repression of tyrosine-phosphorylation of the extracellular signal-regulated kinase 2 selectively defines the two phenotypes. *Mol Biol Cell* 1994; 5: 497-509.

Melton DA. Translocation of a localized maternal mRNA to the vegetal pole of *Xenopus*

oocytes. *Nature* 1987; 328: 80-82.

Melton DA. Pattern formation during animal development. *Science* 1991; 252: 234-241.

Mercer RW, Biemesderfer D, Bliss DP, Collins JH, Forbush B. Molecular cloning and immunological characterization of the γ -polypeptide, a small protein associated with Na, K-ATPase. *J Cell Biol* 1993; 121: 579-586.

Millen KJ, Wurst W, Herrup K, Joyner AL. Abnormal embryonic cerebellar development and patterning of postnatal foliation in two mouse Engrailed-2 mutants. *Development* 1994; 120: 695-706.

Millet S, Alvarado-Mallart RM. Expression of homeobox-containing gene En-2 during the development of the chick central nervous system. *Eur J Neuroscience*. 1995; 7: 777-791.

Miquel J. An update on mitochondrial-DNA mutation hypothesis of cell aging. *Mutation Res* 1992; 275: 209-216.

Minturn JE, Geschwind DH, Fryer HJL, Hockfield S. Early postmitotic neurons transiently express TOAD-64, a neural specific protein. *J Comp Neurol* 1995; 355: 369-379.

Mohandas T, Sparkes RS, Suh EJ, Eersfield MS. Regional localization of the human genes for S-adenosyl-homocysteine hydrolase (cen \rightarrow Q13) and adenosine deaminase (Q13 \rightarrow qter) on chromosome 20. *Hum Genet* 1984; 66: 292-295.

Moment GB. Theories of aging: An overview. In: *Testing the Theories of Aging*, Adelman RC, Roth GS, eds. Boca Raton, FL, CRC, 1982, pp 1-23.

Monuki ES, Weinmaster G, Kuhm R, Lemke G. SCIP: A glial POU domain gene regulated by cyclic AMP. *Neuron* 1989; 3: 783-793.

Moorman JR, Ackerman SJ, Kowdley GC, Griffin MP, Mounsey JP, Chen Z, Cala SE, O'Brain JJ, Szabo G, Jones LR. Unitary anion currents through phospholemman channel molecules. *Nature* 1995; 377: 737-740.

Moorman JR, Palmer CJ, John JE, Durieux ME, Jones LR. Phospholemman expression induces a hyperpolarization-activated chloride current in *Xenopus* oocytes. *J Biol Chem* 1992; 267: 14551-14554.

Mon N, Stein R, Sigmund O, Anderson DJ. A cell type-specific silencer element that controls the neural-specific expression of the SCG10 gene. *Neuron* 1990; 4: 583-594.

Morris CM, Honeybone LM, Hollings PE, Fitzgerald PH. Localization of the SRC oncogene to chromosome band 20q11.2 and loss of this gene with deletion (20q) in two leukemic patients.

Blood 1989; 74: 1768-1773.

Murphy P, Davidson DR, Hill RE. Segment-specific expression of a homeobox-containing gene in the mouse hindbrain. *Nature* 1989; 341: 156-159.

Nagele RG, Lee HY. Studies on the mechanism of neurulation in the chick: Microfilament-mediated changes in cell shape during uplifting of neural folds. *J Exp Zool* 1980; 213: 391-398.

Nagele RU, Lee HY. Studies on the mechanism of neurulation in the chick: Morphometric analysis of the relationship between regional variations in cell shape and sites of motive force generation. *J Exp Biol* 1987; 24: 197-205.

Navarre C, Catty P, Leterme S, Dietrich F, Goffeau A. Two distinct genes encode small isoproteolipids affecting plasma membrane H⁺-ATPase activity of *Saccharomyces cerevisiae*. *J Biol Chem* 1994; 269: 21262-21268.

Navarre C, Ferroud C, Ghislain M, Goffeau A. A proteolipid associated with the plasma membrane H⁺-ATPase of fungi. *Ann NY Acad Sci* 1992a; 671: 189-194.

Navarre C, Ghislain M, Leterme S, Ferroud C, Dufour J P, Goffeau A. Purification and complete sequence of a small proteolipid associated with the plasma membrane H⁺-ATPase of *Saccharomyces cerevisiae*. *J Biol Chem* 1992b; 267: 6425-6428.

Nichols DH. Neural crest formation in the head of the mouse embryo as observed using a new histological technique. *J Embryol Exp Morphol* 1981; 64: 105-120.

Nieto MA, Gilardi-Hebenstreit P, Charnay P, Wilkinson DG. A receptor protein kinase implicated in the segmental patterning of the hindbrain and mesoderm. *Development* 1992; 116: 1137-1150.

Nieuwkoop PD. The formation of the mesoderm in urodele amphibians. I. Induction by the endoderm. *Wilhelm Roux Arch Entwicklungsmech Org* 1969; 162: 341-373.

Nieuwkoop PD. The "organization center" of the amphibian embryo: Its origin, spatial organization and morphogenetic action. *Adv Morphogen* 1973; 10: 1-39.

Ning Y, Shay JW, Lovell M, Taylor L, Ledbetter DH, Pereira-Smith OM. Tumor suppression by chromosome 11 is not due to cellular senescence. *Exp Cell Res* 1991a; 192: 220-226.

Ning Y, Weber JL, Killary AM, Ledbetter DH, Smith JR, Pereira-Smith OM. Genetic analysis of indefinite division in human cells: Evidence for a cell senescence-related gene(s) on chromosome 4. *Proc Natl Acad Sci USA* 1991b; 88: 5635-5659.

Nottebohm F, O'Loughlin B, Gould K, Yohay K, Alvarez-Buylla A. The life span of new

neurons in a song control nucleus of the adult canary brain depends on the time of year when these cells are born. *Proc Natl Acad Sci USA* 1994; 91: 7849-7853.

Nusse R, Varmus HE. *Wnt* Genes. *Cell* 1992; 69: 1073-1087.

O'Reilly MA, Smith JC, Cunliffe V. Patterning of the mesoderm in *Xenopus*: Dose-dependent and synergistic effects of Brachyury and Pintallavis. *Development* 1995; 121: 1351-1359.

Orgel LE. The maintenance of the accuracy of protein synthesis and its relevance to ageing. *Proc Natl Acad Sci USA* 1963; 49: 517-521.

Orr WC, Sohal RS. Extension of life span by overexpression of superoxide dismutase and catalase in *Drosophila melanogaster*. *Science* 1994; 263: 1128-1130.

Otte AP, Kramer IJM, Durston AJ. Protein kinase C regulation of the local competence of *Xenopus* ectoderm. *Science* 1991; 251: 570-573.

Otte AP, Moon RT. Protein kinase C isozymes have distinct roles in neural induction and competence in *Xenopus*. *Cell* 1992; 11: 115-132.

Otte AP, van Run P, Heideveld M, van Driel R, Durston AJ. Neural induction is mediated by cross-talk between the protein kinase C and cyclic AMP pathways. *Cell* 1989; 58: 641-648.

Palmer CJ, Scott ET, Jones LR. Purification and complete sequence determination of the major plasma membrane substrate for cAMP-dependent protein kinase and protein kinase C in myocardium. *J Biol Chem* 1991; 266: 11126-11130.

Palmer DN, Fearnley IM, Walker JE, Hall NA, Lake BD, Wolfe LS, Haltia M, Martinus RD, Jolly RD. Mitochondrial ATP synthase subunit c storage in the ceroid-lipofuscinoses (Batten Disease). *Am J Med Genet* 1992; 42: 561-567.

Pang JH, Chen KY. Global change of gene expression at late G1/S boundary may occur in human IMR-90 diploid fibroblasts during senescence. *J Cell Physiol* 1994; 160: 531-538.

Partridge L, Barton NH. Optimality, mutation and the evolution of ageing. *Nature* 1993; 362: 305-311.

Partridge L, Farquhar M. Sexual activity reduces life span of male fruit flies. *Nature* 1981; 294: 580-582.

Pathak B, Neumann JC, Croyle ML, Lingrel JB. The presence of both negative and positive elements in the 5'-flanking sequence of the rat Na, K-ATPase $\alpha 3$ subunit gene are required for brain expression in transgenic mice. *Nuc Acids Res* 1994; 22: 4748-4755.

- Pennisi F. Premature aging gene discovered. *Science* 1996; 272: 193-194.
- Pereira-Smith OM, Fisher SF, Smith ER. Senescent and quiescent cell inhibitors of DNA synthesis. *Exp Cell Res* 1985; 160: 297-306.
- Pfaff SL, Mendelsohn M, Stewart CL, Edlund T, Jessell TM. Requirement for LIM homeobox gene *Is11* in motor neuron generation reveals a motor neuron-dependent step in interneuron differentiation. *Cell* 1996; 84: 309-320.
- Pienta KJ, Getzenberg RH, Coffey DS. Characterization of nuclear morphology and nuclear matrices in ageing human fibroblasts. *Mech Ageing Dev* 1992; 62: 13-24.
- Placzek M, Jessell TM, Dodd J. Induction of floor plate differentiation by contact-dependent, homeogenetic signals. *Development* 1993; 117: 205-218.
- Porteus MH, Bulfone A, Ciaranello RD, Rubenstein JL. Isolation and characterization of a novel cDNA clone encoding a homeodomain that is developmentally regulated in the ventral forebrain. *Neuron* 1991; 7: 221-229.
- Price J, Thurlow L. Cell lineage in the rat cerebral cortex: A study using retroviral-mediated gene transfer. *Development* 1988; 104: 473-482.
- Price M, Lemaistre M, Pischetola M, DiLauro R, Duboule D. A mouse gene related to *Distal-less* shows restricted expression in the developing forebrain. *Nature* 1991; 351: 748-751.
- Price M, Lazzaro D, Pohl T, Mattel MG, Ruther U, Olivo JC, Duboule D, DiLauro R. Regional expression of homeobox gene *Nkx-2.2* in the developing mammalian forebrain. *Neuron* 1992; 8: 241-255.
- Price SR, Nightingale MS, Bobak DA, Tsuchiya M, Moss J, Vaughan M. Conservation of a 23-kDa human transplantation antigen in mammalian species. *Genomics* 1992; 14: 959-964.
- Prowse KR, Greider CW. Developmental and tissue-specific regulation of mouse telomerase and telomere length. *Proc Natl Acad Sci USA* 1995; 92: 4818-4822.
- Quintrell N, Lebo R, Vermus H, Bishop JM, Pettenati MJ, LeBeau MM, Diaz MO, Rowley JD. Identification of a human gene (HCK) that encodes a protein-tyrosine kinase and is expressed in hemopoietic cell. *Mol Cell Biol* 1987; 7: 2267-2275.
- Raff MC, Barres BA, Burne JF, Coles HS, Ishizaki Y, Jacobson MD. Programmed cell death and the control of cell survival: Lessons from the nervous system. *Science* 1993; 262: 695-700.
- Rakic P. Mode of cell migration to the superficial layers of fetal monkey neocortex. *J Comp Neurol* 1972; 145: 61-84.

- Rao KM, Cohen HJ. Actin cytoskeletal network in aging and cancer. *Mutation Res* 1991; 256: 139-148.
- Rattan SIS. DNA damage and repair during cellular aging. *Jut Rev Cytol* 1989; 116: 47-88.
- Riabowol K, Schiff J, Oilman MZ. Transcription factor AP-1 activity is required for initiation of DNA synthesis and is lost during cellular aging. *Proc Natl Acad Sci USA* 1992; 89: 157-161.
- Richardson A, Cheung HT. The relationship between age-related changes in gene expression, protein turnover, and the responsiveness of an organism to stimuli. *Life Sci* 1982; 31: 605-613.
- Roelink H, Augsburger A, Heemskerk J, Korzh V, Norlin S, Ruiz i Altaba A, Tanabe Y, Placzek M, Edlund T, Jessell TM, Dodd J. Floor plate and motor neuron induction by vhh-1, a vertebrate homolog of hedgehog expressed by the notochord. *Cell* 1994; 76: 761-775.
- Romanes GJ. In: *Darwin and After Darwin*. Open Court Publishing Co., London, UK, 1901.
- Rosenberger RF. The initiation of senescence and its relationship to embryonic cell differentiation. *Bioessays* 1995; 17: 257-260.
- Ruiz JC, Robertson EJ. The expression of the receptor-protein tyrosine kinase gene, *eck*, is highly restricted during early mouse development. *Mech Dev* 1994; 46: 87-100.
- Sager R. Senescence as a mode of tumor suppression. *Environ Health Perspect* 1991; 93: 59-62.
- Sager R, Anisowicz A, Neveu M, Liang P, Sotiropoulou G. Identification by differential display of alpha 6 integrin as a candidate tumor suppressor gene. *FASEB J* 1993; 7: 964-970.
- Sambrook J, Fritsch EF, Maniatis T. In: *Molecular Cloning. A Laboratory Manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Plainview, NY 1989, 125-128.
- Sandhu AK, Hubbard K, Kaur GP, Jha KK, Ozer HL, Athwal RS. Senescence of immortal human fibroblasts by the introduction of normal human chromosome 6. *Proc Natl Acad Sci USA* 1994; 91: 5498-5502.
- Sanger F, Milden S, Coulson AR. DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci USA* 1977; 74: 5463-5467.
- Sapolsky RM. In: *Stress, the Aging Brain, and the Mechanisms of Neuron Death*. MIT Press, Cambridge, Mass, 1992, pp 30.
- Sargent TD. Isolation of differentially expressed genes. *Meth Enzymol* 1987; 152: 423-432.

Sasai Y, Lu B, Steinbeisser H, Geissert D, Gont LK, De Robertis EM. *Xenopus chordin*: A novel dorsalizing factor activated by organizer-specific homeobox genes. *Cell* 1994; 79: 779-790.

Sauer FC. Mitosis in the neural tube. *J Comp Neurol* 1935; 62: 277-405.

Saunders NA, Smith RJ, Jetten AM. Regulation of proliferation-specific and differentiation-specific genes during senescence of human epidermal keratinocyte and mammary epithelium cells. *Biochem Biophys Res Commun* 1993; 197: 46-54.

Scheibel ME, Lindsay RD, Tomiyasu U, Scheibel AB. Progressive dendritic changes in aging human cortex. *Exp Neurol* 1975; 47: 392-403.

Schneider-Manoury S, Topilko P, Seitanidou T, Levi G, Cohen-Tannoudji M, Pournin S, Babinet C, Charnay P. Disruption of *Krox-20* results in alteration of rhombomeres 3 and 5 in the developing hindbrain. *Cell* 1993; 79: 1199-1214.

Schoenwolf GC. Cell movements driving neurulation in avian embryos. *Development* 1991; 2 (Suppl): 157-168.

Schoenwolf GC, Desmond NE. Descriptive studies of the occlusion and reopening of the spinal canal of the early chick embryo. *Anat Rec* 1984; 209: 251-263.

Schulte-Merker S, Smith JC. Mesoderm formation in response to Brachyury requires FGF signalling. *Curr Biol* 1995; 5: 62-67.

Schulte-Merker S, Smith JC, Dale L. Effects of truncated activin and FGF receptors and of follistatin on the inducing activities of BVg1 and activin: Does activin play a role in mesoderm induction? *EMBO J* 1994; 13: 3533-3541.

Scott MP. Vertebrate homeobox gene nomenclature. *Cell* 1992; 71: 55-553.

Seshadri T, Campisi J. Repression of c-fos transcription and an altered genetic program in senescent human fibroblasts. *Science* 1990; 247: 205-209.

Seshadri T, Uzman JA, Oshima J, Campisi J. Identification of a transcript that is down-regulated in senescent human fibroblasts. Cloning, sequence analysis, and regulation of the human L7 ribosomal protein gene. *J Biol Chem* 1993; 268: 18474-18480.

Sham MH, Vesque C, Nonchev S, Marshall H, Frain M, Das Gupta R, Whiting J, Wilkinson D, Charnay P, Krumlauf R. The zinc finger gene *Krox20* regulates *HoxB2* (*Hox2.8*) during hindbrain segmentation. *Cell* 1993; 72: 183-196.

Sharpe CR, Fritz A, De Robertis EM, Gurdon JB. A homeobox-containing marker of posterior

neural differentiation shows the importance of predetermination in neural induction. *Cell* 1987; 50: 749-758.

Shibanuma M, Mashimo J, Kuroki T, Nose K. Characterization of the TGF- β 1-inducible hic-5 gene that encodes a putative novel zinc finger protein and its possible involvement in cellular senescence. *J Biol Chem* 1994; 269: 26767-26774.

Sibille C, Chomez P, Wildmann C, Van Pel A, De Plaen E, Maryanski JL, de Bergeyck V, Boon T. Structure of the gene of tuni transplantation antigen P 198: A point mutation generates a new antigenic peptide. *J Exp Med* 1990; 172: 35-45.

Sidman RL, Miale IL, Feder N. Cell proliferation and migration in the primitive ependymal zone; an autoradiographic study of histogenesis in the nervous system. *Exp Neurol* 1959; 1: 322-333.

Simeone A, Acampora D, Gulisano M, Stornaiuolo A, Boncinelli E. Nested expression domains of four homeobox genes in developing rostral brain. *Nature* 1992a; 358: 687-690.

Simeone A, Gilisano M, Acampora D, Stornaiuolo A, Rambaldi M, Boncinelli E. Two vertebrate homeobox genes related to the *Drosophila empty spiracles* gene are expressed in the embryonic cerebral cortex. *EMBO J* 1992b; 11: 135-137.

Smeal T, Claus J, Kennedy B, Cole F, Guarente L. Loss of transcriptional silencing causes sterility in old mother cells of *S. cerevisiae*. *Cell* 1996; 84: 633-642.

Smith JC. Mesoderm-inducing factors in early vertebrate development. *EMBO J* 1993; 12: 4463-4470.

Smith JC, Price BMJ, van Nimmen K, Huylebroeck D. Identification of a potent *Xenopus* mesoderm-inducing factor as a homologue of activin A. *Nature* 1990; 345: 729-731.

Smith JL, Schoenwolf G. Notochordal induction of cell wedging in the chick neural plate and its role in neural tube formation. *J Exp Zool* 1989; 250: 49-62.

Smith JL, Schoenwolf GC. Further evidence of extrinsic forces in bending of the neural plate. *J Comp Neurol* 1991; 307: 225-236.

Smith JR, Pereira-Smith OM. Replicative senescence: Implications for *in vivo* aging and tumor suppression. *Science* 1996; 273: 63-67.

Smith WC, Harland, RM. Injected *wnt-8* RNA acts early in *Xenopus* embryos to promote formation of a vegetal dorsalizing center. *Cell* 1991; 67: 753-765.

Smith WC, Harland, RM. Expression cloning of *noggin*, a new dorsalizing factor localized to

the Spemann organizer in *Xenopus* embryos. *Cell* 1992; 70: 829-840.

Smith WC, Knecht AK, Wu M, Harland RM. Secreted noggin mimics the Spemann organizer in dorsalizing *Xenopus* mesoderm. *Nature* 1993; 361: 547-549.

Smith WC, McKendry R, Ribisi S, Harland RM. A nodal-related gene defines a physical and functional domain within the Spemann organizer. *Cell* 1995; 82: 37-46.

Sohal RS, Agarwal A, Agarwal S, Orr WC. Simultaneous overexpression of copper- and zinc-containing superoxide dismutase and catalase retards age-related oxidative damage and increases metabolic potential in *Drosophila melanogaster*. *J Biol Chem* 1995; 270: 15671-15674.

Sohal RS, Brunk UT. Mitochondrial production of pro-oxidants and cellular senescence. *Mutation Res* 1992; 275: 295-304.

Sohal RS, Weindruch R. Oxidative stress, caloric restriction, and aging. *Science* 1996; 273: 59-63.

Spemann H. In: *Embryonic Development and Induction*. Yale University Press, New Haven, 1938.

Spemann H, Mangold H. Induction von embryonalanlagen durch implantation artfremder organisatoren. *Arch Mikrosk Anat Entwicklungsmech* 1924; 100: 599-638.

Stein GH, Drullinger LF, Robortye RS, Pereira-Smith OM, Smith JR. Senescent cells fail to express cdc2, cycA, and cycB in response to mitogen stimulation. *Proc Natl Acad Sci USA* 1991; 88: 11012-11016.

Stella AMG, Lajtha A. Macromolecular turnover in brain during aging. *Gerontology* 1987; 33: 136-148.

Stitt TN, Gasser UE, Hatten ME. Molecular mechanisms of glial-guided neuronal migration. *Ann NY Acad Sci* 1991; 633: 113-121.

Stokkermans J, van Dongen W, Kaan A, van den Berg W, Veeger C. hdy, a gene from *Desulfovibrio vulgaris* (Hildenborough) encodes a polypeptide homologous to the periplasmic hydrogenase. *FEMS Microbiol Lett* 1989; 58: 217-222.

Storey K, Crossley JM, De Robertis E, Norris WE, Stern CD. Neural induction and regionalization in the chick embryo. *Development* 1992; 114: 729-741.

Strehler B, Hirsch G, Gussek D, Johnson R, Bick M. Codon-restriction theory of aging and development. *J Theor Biol* 1971; 33: 429-474.

- Suzuki N, Rohdewohld H, Neuman T, Gruss P, Scholer HR. Oct-6: A POU transcription factor expressed in embryonal stem cells and the developing brain. *EMBO J* 1990; 9: 3723-3732.
- Tada M. Molecular structure and function of phospholamban in regulating the calcium pump from sarcoplasmic reticulum. *Ann NY Acad Sci* 1991; 671: 92-103.
- Tanford C. The hydrophobic effect and the organization of living matter. *Science* 1978; 200: 1012-1018.
- Thiel G, Schoch S, Petersohn D. Regulation of synapsin I gene expression by the zinc finger transcription factor zif268/egr-1. *J Biol Chem* 1994; 269: 15294-15301.
- Thomsen GH, Melton DA. Processed Vgl protein is an axial mesoderm inducer in *Xenopus*. *Cell* 1993; 74: 433-441.
- Tilney F. Behavior in its relation to the development of the brain. Part II. Correlation between the development of the brain and behavior in the albino rat from embryonic states to maturity. *Bull Neurol Inst NY* 1933; 2: 252-358.
- Tremblay P, Gruss P. Pax: genes for mice and men. *Pharmac Ther* 1994; 61: 205-226.
- Tsering C. Demonstration of segmental arrangement of thoracic spinal motor neurons using lipophilic dyes. *Dev Neurosci* 1992; 14: 308-311.
- Tsuchida T, Ensini M, Morton SB, Baldassare M, Edlund T, Jessell TM, Pfaff SL. Topographic organization of embryonic motor neurons defined by expression of LIM homeobox genes. *Cell* 1994; 79: 957-970.
- Turner DL, Cepko CL. A common progenitor for neurons and glia persists in rat retina late in development. *Nature* 1987; 328: 131-136.
- Valcarce C, Navarrete RM, Encabo P, Loeches E, Satrustegui J, Cuezva JM. Postnatal development of rat liver mitochondrial functions. *J Biol Chem* 1988; 263: 7767-7775.
- Valtorta F, Benfenati F, Greengard P. Structure and function of the synapsins. *J Biol Chem* 1992; 267: 7195-7198.
- Van Allen MI, Kalousek DK, Chernoff GF, Juriloff D, Harris M, McGillivray BC, Yong S-L, Langlois S, MacLeod PM, Chitayat D, Friedman JM, Wilson RD, Mcfadden D, Pantzar S, Ritchie S, Hall JG. Evidence for multi-site closure of the neural tube in humans. *Am J Med Genet* 1993; 47: 723-743.
- Van Straaten HWM, Hekking JWM, Beurgens JPWM, Terwindt-Rouwerthorst E, Drukker J.

- Effect of the notochord on proliferation and differentiation in the neural tube of the chick embryo. *Development* 1989; 107: 793-539.
- Van Straaten HWM, Hekking JWM, Wiertz-Hoessels EJLM, Thors F, Drukker J. Effect of the notochord on the differentiation of a floor plate area in the neural tube of chick embryo. *Anat Embryol* 1988; 177: 317-324.
- Vaziri H, Schachter F, Uchida I, Wei L, Zhu X, Effros R, Cohen D, Harley CB. Loss of telomere DNA during aging of normal and trisomy 21 human lymphocytes. *Am J Hum Genet* 1993; 52: 661-667.
- Von Baer KE. *Über Entwicklungsgeschichte der Thiere, Beobachtung und Reflexion*. I Theil, 1828; II Theil, 1837. Gebrüder Bornträger, Königsberg, 1828, 1837.
- Von Zglinicki T, Saretzki G, Docke W, Lotze C. Mild hyperoxia shortens telomeres and inhibits proliferation of fibroblasts: A model for senescence? *Exp Cell Res* 1995; 220: 186-193.
- Wada K, Aota S, Tsuchiya R, Ishibashi F, Gojobori T, Ikemura T. Codon usage tabulated from the GenBank genetic sequence data. *Nuc Acids Res* 1990; 18: 2367-2411.
- Waddington CH. Induction by the primitive streak and its derivatives in the chick. *J Exp Biol* 1933; 10: 38-46.
- Wadhwa R, Pereira-Smith OM, Reddel RR, Sugimoto Y, Mitsui Y, Kaul SC. Correlation between complementation group for immortality and the cellular distribution of mortalin. *Exp Cell Res* 1995; 216: 101-106.
- Wahle E, Keller W. The biochemistry of 3'-end cleavage and polyadenylation of messenger RNA precursors. *Ann Rev Biochem* 1992; 61: 419-440.
- Wang E. Rapid disappearance of statin, a nonproliferating and senescent cell-specific protein, upon reentering the process of cell cycling. *J Cell Biol* 1985; 101: 1695-1702.
- Wang E, Lee MJ, Pandey S. Control of fibroblast senescence and activation of programmed cell death. *J Cell Biochem* 1994; 54: 432-439.
- Wang E, Tomaszewski G. Granular presence of terminin is the marker to distinguish between senescent and quiescent states. *J Cell Physiol* 1991; 147: 514-522.
- Wang T, Danielson PD, Li B, Shah PC, Kim SD, Donahoe PK. The p21^{RAS}Farnesyltransferase a subunit in TGF- β and activin signaling. *Science* 1996; 271: 1120-1122.
- Wawrzynow A, Theibert JL, Murphy C, Jona I, Martonosi A, Collins JH. Sarcoplipin, the "proteolipid" of skeletal muscle sarcoplasmic reticulum, is a unique, amphipathic, 31-residue

peptide. *Arch Biochem Biophys* 1992; 298: 620-623.

Weissmann A. The duration of life. In: *Collected Essays on Heredity and Kindred Biological Problems*, Poluton EB, ed. Clarendon, Oxford, UK, 1889.

Welsh J, Chada K, Dalal SS, Cheng R, Ralph D, McClelland M. Arbitrarily primed PCR fingerprinting of RNA. *Nuc Acids Res* 1992; 20: 4965-4970.

Wetts R, Fraser SE. Multipotent precursors can give rise to all major cell types of the frog retina. *Science* 1988; 239: 1142-1145.

White NJ, Nacheva E, Asimakopoulos FA, Bloxham D, Paul B, Green AR. Deletion of chromosome 20q in myelodysplasia can occur in a multipotent precursor of both myeloid cells and B cells. *Blood* 1994; 83: 2809-2816.

Wijnholds J, Chowdhury K, Wehr R, Gruss P. Segment-specific expression of the *neuronatin* gene during early hindbrain development. *Dev Biol* 1995; 171: 73-84.

Wilkinson DG, Bhatt S, Chavrier P, Bravo R, Charnay P. Segment-specific expression of a zinc-finger gene in the developing nervous system of the mouse. *Nature* 1989; 337: 461-464.

Wolpert L. Positional information and the spatial pattern of cellular differentiation. *J Theor Biol* 1969; 25: 1-47.

Wong A, Boutis P, Hekimi S. Mutations in the *clk-1* gene of *Caenorhabditis elegans* affect developmental and behavioral timing. *Genetics* 1995; 139: 1247-1259.

Yakovlev PI. Paraplegia in flexion of cerebral origin. *J Neuropathol Exp Neurol* 1954; 13: 267-296.

Yamada H. Suppression of tumorigenicity and induction of senescence on human endometrial carcinoma cell lines by transfer of normal human chromosomes. *Hokkaido Igaku Zasshi - Hokkaido J Med sci* 1994; 69: 1443-1454.

Yamada T, Pfaff S, Edlund T, Jessell TM. Control of cell pattern in the neural tube: Motor neuron induction by diffusible factors from notochord and floor plate. *Cell* 1993; 73: 673-686.

Yamakuni T, Kuwano R, Odani S, Miki N, Yamaguchi Y, Takahashi Y. Molecular cloning of cDNA to mRNA for a cerebellar spot 35 protein. *J Neurochem* 1987; 48: 1590-1596.

Yu C-E, Oshima J, Fu Y-H, Wijsman EM, Hisama F, Alisch R, Matthews S, Nakura J, Miki T, Ouais S, Martin GM, Mulligan J, Schellenberg GD. Positional cloning of the Werner's syndrome gene. *Science* 1996; 272: 258-262.

Zheng C, Heintz N, Hatten ME. CNS gene encoding astrotactin, which supports neuronal migration along glial fibers. *Science* 1996; 272: 417-419.

Zs-Nagy I, Nagy K. On the role of cross-linking of cellular proteins in aging. *Mech Ageing Dev* 1980; 14: 245-251.