

Gigi Kroll, MD ♦ Zhanna M. Pinkus, MD

Aileen Slivkoff, NP

Warning Signs During Pregnancy

Finding and dealing with problems early can often lower risks for you and your baby. Here are warning signs to recognize:

- Bleeding or spotting from your vagina.
- A gush or leak of water from your vagina.
- Uterine cramping or tightening 6 or more times per hour if you are less than 37 weeks pregnant.
- Vague signs of preterm labor:
 - Menstrual-like cramping.
 - A dull, low backache.
 - Pelvic pressure or heaviness.
 - Intestinal cramping, without diarrhea.
 - An increase or change in the character of vaginal discharge.
 - A general feeling that "something is not right."
- Sharp, non-stop pain in your belly.
- Fever over 100°F
- Nausea or vomiting that won't go away.
- Sudden swelling of your face, hands, or feet.
- Continued bad headache that won't go away after resting and taking Tylenol.
- Blurred vision or spots before your eyes.
- Pain and/or burning when you urinate.
- Contact with someone who has H1N1 flu, measles, chicken pox, whooping cough, or any other illnesses you are concerned about.
- A decrease in your baby's normal movements and activity. Your baby should move at least 10 separate times in a 2-hour period daily after 28 weeks of pregnancy.
 - See "Fetal Kick Count" document to monitor your babies kick counts from home.