



Center for Sexual & Urinary Function

Internal Penile Pump **HOW TO TAKE MEDICATION**

Please note over the counter medications (OTC) need to be purchased on your own and you may not get or need the assistance from a pharmacist to obtain these prescriptions.

BEFORE THE PROCEDURE:

- Please take **CIPRO 500 mg** the morning and evening before the procedure date, the day of the procedure with a sip of water only and the day after the procedure.

Note: High blood pressure medications, heart medications or the CIPRO prescribed by your doctor must be taken with the smallest amount of water possible in the morning of the procedure, even if advised by the Hospital or the Center not to eat or drink anything after 12:00 midnight.

- Wash your penis, scrotum, groin, and area over pubic bone and between the scrotum and rectum with 45 ml of **Hibiclens solution** twice daily starting 2 days before surgery and finishing the morning of surgery (5 washings). You can purchase the solution at your local pharmacy (Walgreens 236 ml of Hibiclens costs \$9.99)

AFTER THE PROCEDURE:

- **Colace** 100 mg (OTC) (Ducosate)

Stool lubricant: With the pain medication, the majority of patients get constipated and in order to prevent this from occurring, we prescribe Colace to be taken 2 tablets 2 times a day. Please start 3 days before procedure if you have a tendency to become constipated.

- **MOM** (OTC) (Milk of Magnesia)

Stool laxative to be taken if the patient has not had a bowel movement after being home for 24 hours. Use as directed.

- **Keflex** 500 mg (Cephalexin Monohydrate)

Antibiotic: It must be taken 4 times a day for a total of 7 days. To be started after the patient is discharged home.

- **Percocet** (Oxycodone HCl with Acetaminophen and/or APAP)

Pain medication for moderate and strong pain: to be taken after the procedure in case of onset of pain 2 tablets every 4-6 hours.

Peter Niemczyk, MD, FACS ☒ 20325 N 51st Ave, Suite 102, Glendale, AZ 85308

☎ (623) 780-2300 or (928) 537-0111, Fax (623) 582-9666