


Bowel Preparation for Colonoscopy With MoviPrep

Colonoscopy is a routine, generally safe procedure used to examine the colon. There are steps that you, the patient, must take before the procedure to ensure the highest level of safety and effectiveness.

- ① Make arrangements to have someone drive you home after the procedure
 - Colonoscopy involves sedation, and you will not be allowed to leave unaccompanied
- ② Follow your physician's instructions regarding medicines to avoid and diet to follow before your procedure
- ③ Finish the entire bowel prep regimen as described on the following page, unless otherwise directed by a physician

MoviPrep is available at most retail pharmacy outlets. If your pharmacy does not have MoviPrep in stock, it can be ordered and delivered within 24 hours.

Information for your pharmacist: NDC 65649-201-75



MoviPrep
(PEG-3350, Sodium Sulfate, Sodium Chloride, Potassium Chloride,
Sodium Ascorbate and Ascorbic Acid for Oral Solution)

www.MoviPrep.com

Important Safety Information about MOVIPREP

Do not take MOVIPREP® (PEG-3350, sodium sulfate, sodium chloride, potassium chloride, sodium ascorbate and ascorbic acid for oral solution) if you have a gastrointestinal (GI) obstruction (a blockage in your bowel), bowel perforation (an opening in the wall of your stomach or intestine), gastric retention (problems with food and fluid emptying from your stomach), ileus (bowel muscle unable to move stomach contents), toxic colitis (damage to the thickness of the intestinal wall) or toxic megacolon (an extremely dilated or distended colon with fever and/or abdominal pain at times), or if you have a known allergy to MOVIPREP or any of its components.

It is important to drink sufficient liquids before, during, and after the use of MOVIPREP. Liquid and blood salt changes can lead to serious side effects including abnormal heartbeats, seizures, and kidney problems. Talk to your doctor before taking MOVIPREP if you take any medications that increase the risk of blood salt abnormalities or if you have known or suspected low blood salt (sodium) level (hyponatremia).

If you have kidney problems or if you take medication that affects kidney function, talk to your doctor before taking MOVIPREP.

If you have suspected bowel blockage or a suspected opening in the wall of your stomach or intestine, talk to your doctor before taking MOVIPREP.

Talk to your doctor before taking MOVIPREP if you have problems swallowing, if you have gastric reflux, or if you are predisposed to aspiration.

If you have a condition that destroys red blood cells called glucose-6-phosphate dehydrogenase (G6PD) deficiency, talk to your doctor before taking MOVIPREP.

Please be advised that MOVIPREP contains 131 mg of phenylalanine per treatment.

Important Safety Information continued on back.


Patient Instructions

The MoviPrep carton contains 4 pouches and a disposable container for mixing. You must complete the entire prep to ensure the most effective cleansing.

Beginning at _____


STEP 1 MIX FIRST DOSE

- Empty 1 Pouch A and 1 Pouch B into the disposable container
- Add lukewarm drinking water to the top line of the container. Mix to dissolve
If preferred, mix solution ahead of time and refrigerate prior to drinking. The reconstituted solution should be used within 24 hours.



STEP 2 DRINK FIRST DOSE

- The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz), until the full liter is consumed
- Drink 16 oz of the clear liquid of your choice. This is a necessary step to ensure adequate hydration and an effective prep




Clear liquids include water, ginger ale, apple juice, Gatorade*, lemonade, and broth. No red or purple liquids. Ask your doctor if you have any questions about whether a particular drink is acceptable.

*Gatorade is a registered trademark of Stokely-Van Camp, Inc., Chicago, IL.

Beginning at _____


STEP 3 MIX SECOND DOSE

- Empty 1 Pouch A and 1 Pouch B into the disposable container
- Add lukewarm drinking water to the top line of the container. Mix to dissolve
If preferred, mix solution ahead of time and refrigerate prior to drinking. The reconstituted solution should be used within 24 hours.



STEP 4 DRINK SECOND DOSE

- The MoviPrep container is divided by 4 marks. Every 15 minutes, drink the solution down to the next mark (approximately 8 oz), until the full liter is consumed
- Drink 16 oz of the clear liquid of your choice. This is a necessary step to ensure adequate hydration and an effective prep



Clear liquids include water, ginger ale, apple juice, Gatorade*, lemonade, and broth. No red or purple liquids. Ask your doctor if you have any questions about whether a particular drink is acceptable.

*Gatorade is a registered trademark of Stokely-Van Camp, Inc., Chicago, IL.

A colonoscopy prep causes the body to lose a significant amount of fluid and can result in sickness due to dehydration. It's important that you prepare your body by drinking extra clear liquids before the prep. Stay hydrated by drinking all required clear liquids during the prep. Replenish your system by drinking clear liquids after returning home from your colonoscopy.

If you have any questions, please call our office at _____.

Important Safety Information about MOVIPREP (cont)

If you are pregnant or planning to become pregnant, talk to your doctor before taking MOVIPREP.

In clinical studies with MOVIPREP, the most common side effects for split dosing were malaise (feeling uneasy), nausea, stomach (abdominal) pain, vomiting, and upper stomach pain. The most common side effects for evening only dosing were stomach swelling (abdominal distension), anal discomfort, thirst, nausea, stomach (abdominal) pain, sleep disorder, rigors (body shakes), hunger, malaise (feeling uneasy), vomiting, and dizziness.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch/ or call 1-800-FDA-1088.

For product information, adverse event reports, and product complaint reports, please contact:

Salix Product Information Call Center
Phone: 1-800-508-0024
Fax: 1-510-595-8183
Email: salix@medcomsol.com

Please see complete Prescribing Information for MOVIPREP at www.MoviPrep.com.



Website: www.salix.com 8510 Colonnade Center Drive, Raleigh, NC 27615 Tel • 866.669.SLXP (7597)
©2013 Salix Pharmaceuticals, Inc. All rights reserved. Printed in USA. MOV 11/43-2

MoviPrep
(PEG-3350, Sodium Sulfate, Sodium Chloride, Potassium Chloride,
Sodium Ascorbate and Ascorbic Acid for Oral Solution)

www.MoviPrep.com