

Jennifer Brenner, MD

Jeffrey Frank, MD

Kevin Rufner, MD

Kevin Sieja, MD

Thomas E. Trouillot, MD

# **SUPREP Bowel Prep**

#### 2 to 3 days before your colonoscopy:

Avoid raw fruits, raw vegetables and grains (especially whole wheat, seeds or nuts) for 2 days before the procedure.

# 1 day before your colonoscopy:

Enjoy a light breakfast such as cereal, toast, eggs (but without seeds, raw fruits or raw vegetables). After your breakfast meal DO NOT eat solid food until after your colonoscopy is performed the following day. YOU MAY HAVE ONLY CLEAR LIQUIDS UP TO TWO (2) HOURS BEFORE YOUR PROCEDURE. Chewing Gum <u>is</u> acceptable. Clear liquids include water, apple juice, ginger ale, 7-up, coffee (black), tea (no milk or creamers), Gatorade (not red in color), jello (not red in color), chicken broth (fat free) or beef broth (fat free).

# The day before your colonoscopy:

#### At 6 PM:

- Step 1: Pour ONE (1) 6-ounce bottle of SUPREP liquid into the mixing container.
- Step 2: Add drinking water to the 16-ounce line on the container and mix.
- Step 3: Drink ALL the liquid in the container.
- Step 4: You must drink two (2) more 16-ounce containers of water over the next hour.

Continue drinking clear liquids until you go to bed.

# The day of your colonoscopy:

FOUR (4) hours before your procedure, repeat Steps 1 through 4 as above using the second 6-ounce bottle of SUPREP.

You must finish drinking the final glass of water at least 2 hours before your colonoscopy.

#### DO NOT HAVE ANYTHING TO DRINK FOR 2 HOURS BEFORE YOUR PROCECURE.

For example, if you are scheduled for a 10 A.M. procedure, stop your clear liquid diet and have nothing by mouth after 8 A.M. the morning of your procedure.

Please call us with any questions at 303-861-0808.