



Post Treatment Instructions

Phototherapy BBL™ (Broadband Light)

The natural extrinsic consequences of aging on your skin include sun damage, freckles, age spots, and redness caused by broken capillaries. Other skin defects arise from intrinsic hormonal factors that create an over production of melanin as in melasma, or hereditary factors that produce more vessels as in rosacea.

Phototherapy involves the reduction of these signs of aging and skin defects using non-invasive pulses of Broadband Light (BBL). If brown spots are the target, BBL penetrates the skin to reach the melanocytes. The particles of the cells left behind will peel or slough off within 7-14 days. If redness is the target, blood vessels in the deeper layers of the skin absorb the light and the heat created by the light damages the vessels. The vessels are shut down and the body continues to absorb the destroyed vessel remnants.

There is no recovery time and low risk of complications with Phototherapy BBL™ treatments.

Based on your skin conditions, your clinician will customize a treatment plan with the recommended number and frequency of treatments to achieve your desired result.

Typically, a range of 2-5 sessions is recommended, performed 2-4 weeks apart until your desired result has been achieved.

The treated area must be treated with care. **BE GENTLE!** Do not scratch or pick at your skin

- Until sensitivity has completely subsided, avoid all of the following:
 - Use of scented lotions or soaps, exfoliant creams (Retin-A, glycolic/salicylic and alpha- hydroxy acids), acne creams or gels, loofa sponges and aggressive scrubbing
 - Hot or cold water - wash with tepid water
 - Shaving
 - Swimming pools and spas with multiple chemicals/chlorine
 - Activities that cause excessive perspiration
- A cold compress or an ice pack can be used to provide comfort if the treated area is especially warm. This is typically only needed within the first 12 hours after the treatment.
- Skin may appear red and swollen and have a mild sunburn sensation. This is a normal reaction. It can last from a few hours to days after treatment. Applying an ice pack for the first 24 hours will help minimize swelling.
- Vitamin E or Aloe Vera applied to the treatment area may provide a soothing effect. Post treatment discomfort is typically minimal but if the area is very uncomfortable, oral pain relievers (i.e. Extra Strength Tylenol or Advil) may be used.
- There may be erythema (redness) and slight edema (swelling) around the treated vessels. Often they are lighter in appearance and look somewhat smudgy or less defined.

- Freckles and sunspots may turn slightly darker initially and then flake off within 7-14 days. This is a desired and normal reaction.
- Makeup may be applied immediately after the treatment as long as skin integrity has not been compromised.
- In the unusual case of crusting of the skin in the treated area, apply an antibiotic ointment twice a day to the affected areas. Do not pick at these areas, as this may result in infection or scarring.
- If the skin is broken or a blister appears, apply an antibiotic ointment and **contact the office immediately**. Keep the affected area moist and avoid direct sunlight.
- Strictly avoid any sun exposure to the treated area for a minimum of 7 days after the procedure. The treated area is more prone to sunburn and pigmentation change. Keep the area covered and use a sun block with SPF 30+ (and preferably a physical barrier ingredient like Zinc). Reapply every 2 hours. Delayed blistering secondary to sun exposure has been noted up to 72 hours post treatment.
- **Throughout your PhotoTherapy BBL™ treatment series, sun exposure should be avoided between treatments and a sunblock should be used on a daily basis.**

Patient signature _____

Date _____ Patient ID# _____ Witness _____