

Michael R. Macdonald, MD

BOARD CERTIFIED FACIAL PLASTIC SURGEON



Aesthetic Surgery & Skin Rejuvenation

415.956.3223

500 Sutter Street, Suite 430
San Francisco, CA 94102

Post Treatment Instructions

BBL™ Acne Treatment

Patient response can vary after BBL 420 nm Acne treatment. Erythema (redness) is usually noted within a few minutes of procedure completion. A slight sunburn sensation in and around treated area is also normal, expected. These reactions subside within 1-4 hours of treatment.

The treated area must be handled with care. **BE GENTLE!** Do not scratch or pick at your skin

- Until sensitivity has completely subsided, avoid all of the following:
 - Use of scented lotions or soaps, exfoliant creams (Retin-A, glycolic/salicylic and alpha- hydroxy acids), acne creams or gels, loofa sponges and aggressive scrubbing
 - Hot or cold water - wash with tepid water
 - Shaving
 - Swimming pools and spas with multiple chemicals/chlorine
 - Activities that cause excessive perspiration
- A cold compress or an ice pack can be used to provide comfort if the treated area is especially warm. This is rarely needed but if so, usually within the first 4 hours after the treatment.
- Makeup can be applied immediately after treatment as long as skin is intact and no sign of blistering.
- Vitamin E or Aloe Vera applied to the treatment area may provide a soothing effect.
- Post treatment discomfort is typically minimal but if the area is very uncomfortable, oral pain relievers (i.e. Extra Strength Tylenol or Advil) may be used.
- If the skin is broken or a blister appears, apply an antibiotic ointment and **contact the office immediately**. Keep the affected area moist and avoid direct sunlight.
- Strictly avoid any sun exposure to the treated area for a minimum of 7 days after the procedure. The treated area is more prone to sunburn and pigmentation change. Keep the area covered and use a sun block with a protection factor of 30+ reapplying the sun block every two hours. Delayed blistering secondary to sun exposure has been noted up to 72 hours post treatment.
- Subsequent treatments are based upon your clinician's recommendation. Typical treatment plan is a series of 6-8 sessions. The series may entail 2 treatments per week for 2-4 weeks, or perhaps 1 treatment every other week for a period of 12-16 weeks until acne clears. The number and frequency of treatments needed depends on the severity of your acne and how well it responds to BBL.

- Clinical studies suggest that at least two BBL 420 nm acne treatments are needed to begin to see results and results have been reported to last up to 6 months.

Patient signature _____

Date _____ Patient ID# _____ Witness _____