



Restylane

SILK

SUNNING
Sipping
SMOOCHING
Smiling

When all your living starts lining up


start Silking

Learn why 96% of RealSelf.com users
say *Restylane*® *Silk* is WORTH IT.

Self-reported survey. Accessed August 8, 2016.

Indication: *Restylane*® *Silk* is indicated for submucosal implantation for lip augmentation and dermal implantation for correction of perioral rhytids in patients over the age of 21.

Please see complete Important Safety Information on back.



Living lines are part of aging.

Everyday life means a lot of stress to the sensitive skin around the mouth. That includes eating and drinking, talking, laughing, and all of the ways we express ourselves. So when it comes to aging, it's no wonder the living lines around the mouth continue to develop and deepen.

It's the end of the line for lips.

Restylane® Silk smoothes those living lines around your mouth and delivers fuller lips with natural-looking results. It's a unique formulation of hyaluronic acid, which occurs naturally in youthful skin. And because it's designed for precision, it allows your specialist to create the subtle enhancement you're looking for.

The reviews of *Silk* are in.*

See what over 2300 people said about their *Silk* treatment.

91% saw natural-looking results

88% were satisfied or very satisfied with their *Silk* treatment

85% would recommend *Silk* to a friend or family member

*Data on file. *Restylane Silk* Satisfaction Survey, Fort Worth, TX: Galderma Laboratories, L.P., 2016. Percentages may not total 100 due to rounding.



In a self-reported survey on **RealSelf.com**, **96%** gave *Silk* a **WORTH IT** rating.

Accessed August 8, 2016.

The most commonly observed side effects are swelling, tenderness, bruising, pain, and redness at the injection site. These are typically mild in severity and resolve in 2-7 days after treatment.

Please see complete Important Safety Information on back.

StartSilking.com

Anne, 55



Before

After (Day 7)

Never stops laughing.

Anne's *Silk* story.

Years of laughing left Anne with lines around her mouth. And while she never wants to stop laughing, she was ready for a smooth lip line.

"My lip lines have actually changed. They're smoother, they're fuller. I feel sexier and more confident."

Before

After (Day 7)



1.5 mL of *Restylane*[®] *Silk* in lips.
0.8 mL of *Restylane Silk* in perioral (lip) lines around the mouth.

Actual *Restylane*[®] *Silk* patient. Individual results may vary.

Watch Anne's full *Silk* story
at StartSilking.com



Carol, 61



Before

After (Day 7)

Always sipping.

Carol's *Silk* story.

Carol says she likes to drink with straws. But all that sipping has left her with lines around her mouth.

"My lips are fuller and softer, and it's so much nicer putting lipstick on."

Before

After (Day 7)



0.5 mL of *Restylane Silk* in lips.
0.5 mL of *Restylane Silk* in perioral (lip) lines around the mouth.

Actual *Restylane*[®] *Silk* patient. Individual results may vary.

After treatment, you might notice some swelling, redness, pain, bruising, or tenderness. This is typically mild in severity and normally lasts less than 18 days in lips.

Please see complete Important Safety Information on back.

StartSilking.com

Want more volume in your lips?

If you want even more definition in your lips, *Restylane-L*® could be right for you. Talk to your specialist about how to get your most beautiful results.

See *Restylane-L* results.

Victoria, 50



Before

After (Day 7)

3 mL of *Restylane-L* in nasolabial folds, marionette lines, and upper lip.

Actual *Restylane-L*® patient. Individual results may vary.

Restylane®

Products in the *Restylane* family contain traces of gram-positive bacterial protein and are contraindicated for patients with allergies to such material or in patients with severe allergies that have required in-hospital treatment. These products should not be used by patients with bleeding disorders or by pregnant or breastfeeding women. *Restylane-L*, *Restylane® Silk* and *Restylane® Lyft with Lidocaine* should not be used by anyone with a known allergy to lidocaine. Products should not be injected anywhere except areas indicated in their Instructions for Use.

Please see complete Important Safety Information on back.



GET CONNECTED TO PRICELESS REWARDS

Sign up at [AspireRewards.com](https://www.aspiregalderma.com) for your complimentary ASPIRE Galderma Rewards membership to get exclusive benefits right away.

JOIN FOR FREE.

You'll receive an instant \$20 treatment certificate when you sign up, plus access to exclusive perks and discounts on treatments you love. You also have the opportunity to start raising your status level for even more rewards.

EARN POINTS.

With ASPIRE Galderma Rewards, using Galderma aesthetic products isn't the only way to earn points. We also reward your loyalty in other unique ways throughout your aesthetic journey.

GET REWARDED.

It's easy to redeem your points for a choice of exclusive rewards, including cash back via check cards, savings on Galderma aesthetic treatments from your aesthetic healthcare professional, and shopping in our retail boutique that features brands you know and love.

Ask your specialist for *Restylane® Silk* by name.

Important Safety Information

Indications: The *Restylane* family of products includes *Restylane®*, *Restylane-L®*, *Restylane® Lyft with Lidocaine* and *Restylane® Silk*. *Restylane*, *Restylane-L*, and *Restylane Lyft with Lidocaine* are indicated for the correction of moderate to severe facial wrinkles and folds, such as nasolabial folds.

Restylane and *Restylane-L* are indicated for mid-to-deep dermal implantation. *Restylane* and *Restylane-L* are also indicated for submucosal implantation for lip augmentation in patients over the age of 21.

Restylane Lyft with Lidocaine is indicated for implantation into the deep dermis to superficial subcutis. *Restylane Lyft with Lidocaine* is also indicated for cheek augmentation and for the correction of age-related midface contour deficiencies in patients over the age of 21.

Restylane Silk is indicated for submucosal implantation for lip augmentation and dermal implantation for correction of perioral rhytids in patients over the age of 21.

Products in the *Restylane* family contain traces of gram-positive bacterial protein and are contraindicated for patients with allergies to such material or in patients with severe allergies that have required in-hospital treatment. These products should not be used by patients with bleeding disorders or by pregnant or breastfeeding women. *Restylane-L*, *Restylane Silk* and *Restylane Lyft with Lidocaine* should not be used by anyone with a known allergy to lidocaine. Products should not be injected anywhere except areas indicated in their Instructions for Use.

Use of products in the *Restylane* family at the site of skin sores, pimples, rashes, hives, cysts, or infection should be postponed until healing is complete. The most commonly observed side effects are swelling, redness, pain, bruising, headache, tenderness, and itching at the injection site. These are typically mild in severity and typically resolve in less than 7 days in nasolabial folds and less than 14 days in lips. Serious but rare side effects include delayed onset infections, recurrence of herpetic eruptions, and superficial necrosis at the injection site. Do not implant into blood vessels. Use with caution in patients recently treated with anticoagulant or platelet inhibitors to avoid bleeding and bruising.

The *Restylane* family of products is available only through a licensed practitioner. Complete Instructions for Use are available at www.RestylaneUSA.com.



© 2016 Galderma Laboratories, L.P.
All trademarks are the property
of their respective owners.
MP-SLK-00002 08/16

Restylane®
SILK