

Root Canal Therapy

Homecare:

Now that you have had root canal therapy, it is important to follow these recommendations to ensure healing.

Caring for a temporary restoration

- If we placed a temporary filling or crown on your tooth, avoid chewing for at least 30 minutes to allow the restoration to harden.
- To keep your temporary restoration in place, avoid eating hard or sticky foods, especially chewing gum.
- Chew on the opposite side of your mouth.
- Do not worry if a small portion of the temporary filling wears away or breaks off, but if the entire filling wears out, or if temporary crown comes off, call us so it can be replaced.

Chewing and eating

- If we used anesthetic during procedure, avoid chewing until numbness has worn off. Your lips, teeth and tongue may be numb for several hours.

Brushing and Flossing

- Brush and floss normally

Medications and discomfort

- If antibiotics were prescribed, continue to take them for the indicated length of time, even if all symptoms and signs of infection are gone.
- To control discomfort, take pain medication before the anesthetic has worn off or as recommended. It is normal to experience some discomfort for several days after a root canal appointment, especially when chewing.
- To further reduce discomfort or swelling, rinse your mouth three times a day with warm salt water. Use about a pinch of salt per glass of warm water.

When to call us

- Call our office if your bite feels uneven
- You have sensitivity or discomfort that increases or continues beyond three or four days
- Your temporary filling or crown comes off
- You have any questions or concerns